

JOURNAL OF ALGEBRA 48, 417–438 (1977)

Standard Subgroups of Type  $L_n(2^n)$ 

GARY M. SEITZ\*

*Department of Mathematics, University of Oregon, Eugene, Oregon 97403**Communicated by Marshall Hall, Jr.*

Received September 22, 1976

## INTRODUCTION

In this paper we prove the following classification theorem.

**THEOREM.** *Let  $G$  be a finite group and let  $A$  be a standard subgroup of  $G$ . Assume that  $|Z(A)|$  is odd,  $C_G(A)$  has cyclic Sylow 2-subgroups, and  $\tilde{A} = A/Z(A) \cong L_n(q)$ ,  $q$  even, where  $n \geq 3$  and  $\tilde{A} \not\cong L_3(2), L_4(2)$ . In addition we assume*

*(\*) In core-free sections of  $G$  the 2-layers of centralizers of 2-subgroups are semisimple.*

*Then one of the following holds:*

- (i)  $O(G)A \trianglelefteq G$ ,
- (ii)  $E(G/Z(G)) \cong L_n(q^2)$ .
- (iii)  $E(G/Z(G)) \cong L_n(q) \times L_n(q)$ .

We remark that (\*) is needed only when  $G$  is of type (iii) in order to show that a certain subgroup  $G_0$  with  $G_0/Z(G_0) \cong L_n(q) \times L_n(q)$  is, in fact, normal in  $G$ . Here it is used only when the 2-components involved are of type  $L_n(q)$ . Finally, condition (\*) is a consequence of the  $B$ -conjecture of Thompson.

Combining the above theorem with the work in [1, 6, 7, 18] we have the following.

**COROLLARY.** *Let  $G$  be a finite group with standard subgroup  $A$  satisfying  $|Z(A)|$  odd and  $A/Z(A) \cong L_n(q)$ . In addition assume condition (\*). Then one of the following holds:*

- (i)  $O(G)A \trianglelefteq G$ .
- (ii)  $E(G/Z(G)) \cong L_n(q^2)$ .
- (iii)  $E(G/Z(G)) \cong L_n(q) \times L_n(q)$ .
- (iv)  $E(G/Z(G)) \cong U_3(q)$ ,  $n = 2$ .

\* Supported in part by the NSF under Grant GP 37982X1.

- (v)  $E(G/Z(G)) \cong L_3(q)$ ,  $n = 2$ .
- (vi)  $E(G/Z(G)) \cong J_2$  and  $A \cong L_3(2)$ .
- (vii)  $E(G/Z(G)) \cong HS$ ,  $A_{10}$ , or  $A_{12}$ , and  $A \cong L_4(2)$ .
- (viii)  $E(G/Z(G)) \cong M_{12}$ ,  $A_9$ ,  $J_1$  or  $J_2$ , and  $A \cong L_2(4)$ .
- (ix)  $E(G/Z(G)) \cong G_2(3)$  and  $A \cong L_2(8)$ .
- (x)  $E(G/Z(G)) \cong Sz$  and  $A \cong L_3(4)$ .
- (xi)  $E(G/Z(G)) \cong L_2(25)$ ,  $U_3(5)$ , or  $L_3(5)$ , and  $A \cong L_2(4)$ .

In all cases other than (i), (vii), (viii), and (x),  $C_G(A)$  has Sylow 2-subgroups of order 2. This also occurs in (vii) in  $E(G/Z(G)) \cong HS$  or  $A_{10}$  and in (viii) if  $E(G/Z(G)) \cong M_{12}$  and  $G/O(G) \not\cong \text{Aut}(M_{12})$ . In the remaining cases  $C_G(A)$  has Sylow 2-subgroups as Klein groups.

The case of  $L_n(q)$  has been separated from the work in [16] for the following reason. The induction method of [16] breaks down in low dimensions, so that in any case  $L_3(q)$ ,  $L_4(q)$ , and  $L_5(q)$  would have to be treated separately. However, the alternate method used for the cases  $n \leq 5$  works equally well for any  $n \geq 3$ . The method is to use the parabolic subgroups of  $A$  corresponding to the stabilizer of a nested 1-space and hyperplane of the usual  $n$ -dimensional  $\mathbb{F}_q$ -space for  $SL(n, q)$  in order to construct the corresponding parabolic subgroups of the group  $L_n(q^2)$  or  $L_n(q) \times L_n(q)$ . At this stage it is clear which of (ii) or (iii) holds. For (ii) we use McBride's result [12] to obtain the conclusion. For (iii) we first construct a group  $G_0$  with  $G_0/Z(G_0) \cong L_n(q) \times L_n(q)$  and then use the results of [16, I, Sect. 2]. In independent work Gomi has used similar methods to deal with the case  $A/Z(A) \cong L_3(q)$ ,  $q > 4$ .

Throughout the paper the groups  $A$ ,  $G$  are as in the hypothesis of the theorem. Given a group  $H$  we set  $\tilde{H} = H/Z(H)$ . All groups considered are finite.

## 1. PRELIMINARY LEMMAS

(1.1) Let  $G_0 \cong L_n(q^2)$  with  $n \geq 3$ . Let  $\sigma$  be an involutory field automorphism of  $G_0$  and let  $\tau$  be a graph automorphism of  $G_0$ .

(i)  $O^{2'}(C_{G_0}(\sigma)) \cong L_n(q)$ ,  $O^{2'}(C_{G_0}(\tau)) \cong PSp(n, q^2)$  if  $n$  is even,  $O^{2'}(C_{G_0}(\tau)) \cong PSp(n-1, q^2)$  if  $n$  is odd,  $O^{2'}(C_{G_0}(\sigma\tau)) \cong PSU(n, q)$ .

(ii) If  $n$  is odd, then each involution in  $\text{Aut}(G_0) - G_0$  is conjugate to one of  $\sigma$ ,  $\tau$ ,  $\sigma\tau$ .

(iii) If  $n$  is even, then each involution in  $\text{Aut}(G_0) - G_0$  is conjugate to one of  $\sigma$ ,  $\tau$ ,  $\sigma\tau$ , or  $\tau x$ , where  $x \in G_0$  is a transvection in  $G_0$ . Moreover  $C_{G_0}(\tau x)$  is isomorphic to the centralizer of a transvection in  $PSp(n, q^2)$ .

*Proof.* See [2, Sect. 19].

(1.2) Let  $T$  be a  $p$ -group and  $x$  an element of order  $p$  acting on  $T$ . If  $1 = T_1 < T_1 < \dots < T_n = T$  is a normal series of  $T$  with each  $T_{i+1}/T_i$  elementary and free as an  $\mathbb{F}_p(\langle x \rangle)$ -module, then  $|C_T(x)| = |T|^{1/p}$  and each subgroup of order  $p$  in  $T\langle x \rangle - T$  is conjugate to  $\langle x \rangle$ .

*Proof.* See [20].

(1.3) Let  $G$  be a group and let  $A$  be a standard subgroup of  $G$  with  $C_G(A)$  of 2-rank 1. Let  $S \in \text{Syl}_2(N(A))$ . Assume that  $S \notin \text{Syl}_2(G)$  and  $Z(S) \leq AC_G(A)$ . Then  $[A, O(G)] = 1$  so any 2-component of  $G$  containing  $A$  is necessarily a component of  $G$ .

*Proof.* Given  $G, A, S$  let  $t \in S \cap C_G(A)$  with  $t$  an involution. Choose  $g \in N_G(S) - S$  with  $g^2 \in S$ . Then  $t^g \in Z(S)$  but  $t^g \neq t$ . Let  $tt^g = a$  and  $Y = \langle t, t^g \rangle$ . Write  $X = O(G)$  and  $X = C_X(t)C_X(ta)C_X(a)$ .

Now  $C_X(t) \leq N(A)$ , so  $[A, C_X(t)] \leq A \cap X$  and  $[A, C_X(t)] = 1$ . In particular  $C_X(t) \leq C_X(Y)$ . As  $Y = Y^g$  we also have  $C_X(t^g) \leq C_X(Y)$ . We conclude that  $a \in C_G(X) = C_G(O(G))$ , so  $C_G(O(G)) \geq \langle a^G \rangle \geq A$ .

## 2. LOCAL SUBGROUPS

In this section we determine the structure of certain local subgroups of  $G$ . We assume that  $A$  is a standard subgroup of  $G$  with  $|Z(A)|$  odd and  $\tilde{A} = A/Z(A) \cong L_n(q)$ . We assume that  $G$  is a minimal counterexample to the main theorem. Then  $AO(G) \not\leq G$ . Recall that  $\tilde{A} \cong L_3(2)$  or  $L_4(2)$ .

Let  $K = C_G(A)$ ,  $R \in \text{Syl}_2(K)$ , and  $V \in \text{Syl}_2(A)$ . We are assuming  $R$  is cyclic, so let  $\langle t \rangle = \Omega_1(R)$  and write  $U = Z(V)$ . Then  $U$  is elementary Abelian of order  $q$ . Finally let  $S \in \text{Syl}_2(N(A))$  with  $VR \leq S$ .

- (2.1) (i)  $S \notin \text{Syl}_2(G)$ .
- (ii)  $R = \langle t \rangle$  and  $Ut \subseteq t^G$ .
- (iii) If  $t \neq t^g \in Z(S)$ , then  $V \leq A^g$ .
- (iv)  $t^G \cap U = \emptyset$ .

*Proof.* We first claim that  $Ut \subseteq t^G$ . As  $A$  is transitive on its transvections we need show only that there is some  $t^g \in A^{\#}t$  with  $t^g$  projecting to a transvection in  $A$ . If  $\tilde{A} \cong L_3(q)$  this follows from [16, I, (1.2)] and the fact that  $A$  has just one class of involutions. For  $n \geq 4$  let  $X$  and  $D_{00}$  be as in [16, I, (1.6)]. Then  $\tilde{D}_{00} \cong L_{n-2}(q)$  is simple and [16, I, (1.4)] gives  $t^G \cap A\langle t \rangle \cap C(X) \neq \emptyset$ . Say  $n = 4$ . Then let  $t \neq t^g \in A\langle t \rangle \cap C(X)$ ; so  $t^g$  projects to a transvection in  $A$ . If  $t^g \in A$ , then  $t^g \in C_A(t^g)^{(\infty)}$ , whereas  $t \notin C_G(t)^{(\infty)}$ . So we are done if  $n = 4$ . Now suppose  $n \geq 5$ , set  $Y = O(C_G(X))$ , and let bars denote images in  $C_G(X)/Y$ . By [16, I, (1.10)]  $\overline{D_{00}}$  is standard but not normal in  $\overline{C_G(X)}$ . By the minimality of  $G$  and the corollary we know the structure of  $E(\overline{C_G(X)})$ . In all but one case

it is easy to check that the claim holds. The exceptional case is where  $\overline{D_{00}} \cong L_4(2) \cong A_8$  and  $E(C_G(\overline{X})) \cong A_{10}$ . Here  $t$  acts on  $E(C_G(\overline{X}))$  as a transposition. Identify  $t$  with  $(1, 2)$  and  $\overline{D_{00}}$  with the subgroup of  $A_{10}$  fixing points 1 and 2. From [16, I, (1.4)] we have  $t^h \in D_{00}^* \langle t \rangle$  for some  $h \in G$ . Say  $t^h$  projects to  $a \in D_{00}$ . Then  $a$  is either a transvection or has precisely two nontrivial Jordan blocks (viewing  $a \in SL(6, 2)$ ). In either case,  $a \in C_A(a)'$ . Since  $\tilde{A} \cong L_8(2)$  we have  $t \notin C_G(t)'$ , so  $t^G \cap D_{00} = \emptyset$  and  $t^h = ta$ . We are done if  $a$  is a transvection so suppose this is not the case. Viewing  $\bar{a} \in \overline{D_{00}} \cong A_8$  we may identify  $\bar{a}$  with  $(3, 4)(5, 6)$  and  $\bar{t}^h$  with  $(1, 2)(3, 4)(5, 6)$ . Viewing  $C_G(X) \cap C(t^h)$  as a subgroup of  $C(X)$  and then of  $C(t^h)$  we obtain a contradiction. The claim follows.

Choose  $g \in G$  with  $t^g \in Z(S)$ . Since  $S \in \text{Syl}_2(C(t))$  we may assume that  $g \in N(S)$ . Notice that  $V \in \text{Syl}_2(O^2(C(Z(S))))$  so  $g \in N(V)$ . In particular  $V \leq A^g$ , proving (iii). We know that  $N_A(U)$  is transitive on  $U^\#$  so if (iv) were false then it would be possible to choose  $t^g \in U \cap Z(S)$ . But then  $t^g \in V \leq A^g$  implies that  $t \in A$ , whereas  $|Z(A)|$  is odd. So we also have (iv).

Let  $N = N(S)$ . We have seen that  $N \leq N(V)$ , so  $N$  normalizes each of  $Z(V) = U$  and  $C_S(V) = U \times R$ . If  $|R| > 2$ , then  $t$  is the unique involution in  $\Phi(C_S(V))$ , so  $N \leq C(t)$ , which is not the case. This proves (ii).

Consider the action of  $N$  on  $U \times R$ . We know that  $t^G \cap (U \times R) = t^N \cap (U \times R) = Ut$ . As  $N_A(U)$  is transitive on  $U^\#$ ,  $N$  acts on  $Ut$  as a 2-transitive group of degree  $q$  and  $S \in \text{Syl}_2(C_N(t))$ . In particular (i) holds.

(2.2) *Notation.* Let  $\hat{\phantom{x}}$  denote images in  $A/Z(A)$ . Let  $V_1 = O_2(P_1)$  and  $V_2 = O_2(P_2)$ , where  $\hat{P}_1, \hat{P}_2$  are, respectively, the stabilizer of a 1-space, hyperplane, of the natural module for  $L_n(q)$ . We take  $V \leq P_1 \cap P_2$ , so  $V_i \leq V$ . Then  $\widehat{N_A(V_i)} \supseteq \hat{V}_i(\hat{L}_i \circ \hat{H}_i)$ , where  $\hat{L}_i \cong SL(n-1, q)$  and  $\hat{H}_i$  is cyclic of order dividing  $q-1$  and divisible by  $(1/n)(q-1)$ . Also,  $S \cap N_A(V_i) = VR\langle w \rangle$ , where  $w$  induces a field automorphism on  $A$ .

(2.3) *Let  $X$  act faithfully on the elementary Abelian group  $W \times \langle t \rangle$ , normalizing  $W$  and 2-transitive on  $Wt$ . Suppose that  $C_X(t)$  contains a normal subgroup  $K \cong SL(m, q)$ ,  $m \geq 2$  and  $q = 2^a$ , such that  $K$  acts on  $W$  as on the usual module for  $SL(m, q)$ . Then  $X$  contains a normal elementary subgroup of order  $q^m$ .*

*Proof.* Using the result of Ostrom and Wagner [15] we may assume  $m \geq 3$ . The assumption on  $W$  and  $K$  immediately implies that  $|W| = q^m$  and  $|X : C_X(t)| = q^m$ . Let  $S \in \text{Syl}_2(C_X(t))$  and  $S < \bar{S} \in \text{Syl}_2(X)$ . Then  $X = \bar{S}C_X(t)$ . Choose  $w \in W^\#$  with  $\bar{S} \leq C(w)$  and let  $W_0$  be the corresponding 1-space of  $W$  containing  $w$ , when  $W$  is viewed as the usual module for  $K$ . Then  $S \cap K = C_S(W_0)$  and  $W_0 = C_W(S \cap K)$ .

Consider the group  $Y = N_X(W_0 \langle t \rangle)$ . We know that  $Y \cap K$  induces a solvable group on  $W_0$  of order  $(q-1)b$  with  $b | a$  and is transitive on  $W_0^\#$ . So either  $Y \leq C(t)$  or  $Y$  is 2-transitive on  $W_0 t$  of degree  $q$ . In the latter case look at the group  $C_Y(C(W_0 \langle t \rangle)/O_2 C(W_0 \langle t \rangle)) = Y_0$ . It is easy to see that  $Y_0 \cap C(t)$  is

regular on  $W_0t - \{t\}$ , so  $Y_0$  must induce a Frobenius group of order  $q(q-1)$  on  $W_0\langle t \rangle$ . As  $C_K(W_0\langle t \rangle)$  covers  $C(W_0\langle t \rangle)/O_2(C(W_0\langle t \rangle))$  and is irreducible on  $W/W_0$ , we must have  $Y \leq (Y \cap C(t))C(W/W_0)$ .

We claim that  $N_X(W_0)$  contains a Sylow 2-subgroup of  $X$ . We may assume that  $\bar{S} \cap N_X(W_0) \in \text{Syl}_2(N_X(W_0))$ . First suppose  $\bar{S} \cap N_X(W_0) = \bar{S} \cap Y$ . By the above it follows that  $\bar{S} \cap Y = (\bar{S} \cap C(W_0))\langle j \rangle$ , where  $j \in C_X(t)$  and induces a field automorphism of  $K$ . Say  $g \in N_S(\bar{S} \cap Y) - (\bar{S} \cap Y)$ . Let  $H = \bar{S} \cap K$ . Then  $W_0 = C_W(H) = C_W(\bar{S} \cap C(W_0))$ , so  $H^g \not\leq \bar{S} \cap C(W_0)$ . But then  $\langle j \rangle \neq 1$ ,  $q \geq 4$ , and  $H^g = \Omega_1(H^2)$  has a subgroup of index 2 centralizing  $W_0$ . This forces  $W_0 = C_W(H^g)$  and  $g \in N(W_0)$ , a contradiction. So we have the claim in this case. Now suppose  $\bar{S} \cap N_X(W_0) > \bar{S} \cap Y$ . Then we use the above paragraph in order to apply induction to the group  $N_X(W_0)$  acting on  $W \times \langle t \rangle / W_0 = W/W_0 \times \langle t \rangle W_0/W_0$ . As  $N_K(W_0)$  is transitive on  $(W/W_0)^\#$  and since we are assuming that  $N_X(W_0) \not\leq N(W_0\langle t \rangle)$ , we have a 2-transitive group on  $(W/W_0)tW_0$ . Inductively this group is 2-transitive of degree  $q^{m-1}$  and contains a regular normal subgroup. But then  $\bar{S} \cap N_X(W_0) = (\bar{S} \cap C(W_0))\langle j \rangle$  for  $j \in C_X(t)$  a field automorphism of  $K$ , and we argue as above to get  $\bar{S} = N_S(\bar{S} \cap N(W_0))$ . This proves the claim.

To complete the proof of the lemma we note that  $X = \bar{S}C_X(t) = N(W_0)C_X(t)$  implies that  $X$  is 2-transitive on  $W_0^X = W_0^{C_X(t)}$ . Write  $N_X(W_0) = \bar{S}(N_X(W_0) \cap C_X(t))$ . Then intersecting the conjugates of  $\bar{S}$  in  $N_X(W_0)$  we have  $O_2(N_X(W_0)) \neq 1$ . So by the results of O'Nan [13] and Shult [17] we get the lemma.

(2.4)  $N_G(V_i\langle t \rangle)$  normalizes  $V_i$  and acts on  $tV_i$  as a 2-transitive group of degree  $q^{n-1}$  containing a regular normal subgroup.

*Proof.* This follows immediately from (2.3) once we show that  $X_i = N_G(V_i\langle t \rangle)$  normalizes  $V_i$  and is 2-transitive on  $tV_i$ .  $V_i^\#$  is fused in  $A$ , so by (2.1)(iv)  $t^G \cap V_i = \emptyset$ . Suppose we have  $X_i \not\leq C(t)$ . Then as  $N_A(V_i)$  is transitive on  $V_i^\#$ ,  $t^G \cap V_i\langle t \rangle = tV_i$ ,  $V_i = V_i\langle t \rangle - t^G \leq X_i$ , and  $X_i$  is 2-transitive on  $tV_i$ . So we need only show  $X_i \not\leq C(t)$ .

Let  $t^g \in Ut - \{t\}$ , so by (2.1)(iv),  $V \leq A^g$ . If  $n = 3$  then  $V_1, V_2$  are uniquely determined as the elementary subgroups of  $V$  of maximal order. So  $V_i$  is  $A^g$ -conjugate to  $V_1^g$  or  $V_2^g$ , and so  $N_{A^g}(V_i) \not\leq C(t)$ . If  $n \geq 4$ , then  $V_1V_2 = O_2(C_A(t^g)) \leq (C_A(t^g)^{(\infty)}) < C(t^g)^{(\infty)} \leq A^g$ . Again  $V_1, V_2$  are uniquely determined and we can argue as before.

(2.5) Let bars denote images in  $N_G(V_i\langle t \rangle)/O(N_G(V_i\langle t \rangle))$ . Then  $\overline{N_G(V_i\langle t \rangle)} = \overline{W_i\bar{L}_i\langle i, \bar{w} \rangle}$  where  $W_i > V_i$  is an  $L_i$ -invariant elementary Abelian 2-group,  $W_i/V_i$  is elementary of order  $q^{n-1}$ , and  $W_i/V_i \cong V_i$  as  $L_i$ -modules.

*Proof.* By (2.4)  $\overline{N_G(V_i\langle t \rangle)}$  contains a normal 2-subgroup  $\bar{X}$ . We choose  $X$  a 2-group with  $V_i\langle t \rangle < X$ .  $N_A(V_i)$  contains a cyclic subgroup  $Y_0$  with  $Y_0$  irreducible on  $V_i^\#$ . The map  $\bar{x} \rightarrow [\bar{x}, t]$  gives an  $N_A(V_i)$ -isomorphism between

$\bar{X}/\bar{V}_i\langle t \rangle$  and  $\bar{V}_i$ . Therefore  $\widehat{N_A(V_i)}$  is transitive on  $(\bar{X}/\bar{V}_i\langle t \rangle)^\#$  and it follows that  $\bar{X}/\bar{V}_i$  is elementary Abelian. Let  $\bar{W}_i = [\bar{X}, Y_0]$ . Then  $Y_0$  acts irreducibly on  $\bar{W}_i/\bar{V}_i$  and on  $\bar{V}_i$ .

We claim that  $\bar{W}_i$  is Abelian. Easily  $\bar{V}_i \leq Z(\bar{W}_i)$ . In many cases  $\widehat{N_A(V_i)}$  contains a subgroup  $Y \geq Y_0$  cyclic of order  $q^{n-1} - 1$ . In such cases  $Y$  stabilizes  $\bar{W}_i = [\bar{X}, Y_0]$ . Moreover the map  $\bar{w}_i \rightarrow [\bar{w}_i, t]$  is a  $Y$ -homomorphism from  $\bar{W}_i$  to  $\bar{V}_i$  with kernel  $\bar{V}_i$ . So the claim follows from [6, (2.2)]. Suppose now that no such subgroup,  $Y$ , exists. In particular  $q > 2$  and  $n$  divides  $q - 1$ ; so  $n = 3$  or  $n \geq 5$ .

Let  $\bar{Z}/\bar{V}_i\langle t \rangle$  be a  $GF(q)$ -hyperplane in  $\bar{X}/\bar{V}_i\langle t \rangle$ . Then  $\bar{L}_i$  contains a subgroup  $D$  cyclic of order  $q^{n-2} - 1$  with  $D$  regular on  $(\bar{Z}/\bar{V}_i\langle t \rangle)^\#$  and transitive on  $(\bar{X}/\bar{Z})^\#$ . We write  $\bar{V}_i = \bar{Z}_0 \times \bar{Z}_1$  where  $\bar{Z}_0 = [\bar{Z}, t]$  and  $\bar{Z}_1$  is  $D$ -invariant. Then  $\bar{Z}_0$  is a hyperplane in  $\bar{V}_i$ ,  $\bar{Z}_1$  is a 1-space, and  $D$  is transitive on each of  $\bar{Z}_0$  and  $\bar{Z}_1$ . Next write  $\bar{Z}/\bar{V}_i = \bar{C}/\bar{V}_i \times \langle t \rangle \bar{V}_i/\bar{V}_i$  with  $\bar{C}$   $D$ -invariant, and consider  $\bar{C}/\bar{Z}_0$  and  $\bar{C}/\bar{Z}_1$ .  $\bar{C}/\bar{Z}_1$  is Abelian by [6, (2.2)]. Suppose  $n \geq 5$ . To see that  $\bar{C}/\bar{Z}_0$  is Abelian notice that  $|\bar{C}/\bar{V}_i| = q^{n-2} \geq q^3$ . As  $n \geq 5$   $\bar{C}/\bar{Z}_0$  cannot be a Suzuki 2-group [8], so as  $D$  is transitive on  $(\bar{C}/\bar{V}_i)^\#$ ,  $\bar{C}/\bar{Z}_0$  is actually elementary Abelian. In all cases each of  $\bar{C}/\bar{Z}_1$  and  $\bar{C}/\bar{Z}_0$  are Abelian, which yields  $\bar{C}$  Abelian. By order considerations we see that  $\bar{C}$  is uniquely determined by  $\bar{Z}$ .

Choose  $g \in L_i$  with  $\langle \bar{C}, \bar{C}^g \rangle$  covering  $\bar{X}/\bar{V}_i\langle t \rangle$ . Let  $\hat{W}_i = \langle \bar{C}, \bar{C}^g \rangle$ . Then  $\hat{W}_i$  centralizes  $\bar{C} \cap \bar{C}^g$  and  $\hat{W}_i$  has index of at most 2 in  $\bar{X}$ . Moreover by order considerations  $\bar{C} \cap \bar{C}^g > \bar{V}_i$ . Since  $Z(\bar{X}) = \bar{V}_i$  we have  $\hat{W}_i$  of index 2 in  $\bar{X}$ . If  $h \in N_A(V_i)$  then we again use order considerations to argue that  $\bar{C}^g \cap \bar{C} \cap \bar{C}^h > \bar{V}_i$ . So the above arguments imply that  $\langle \bar{C}, \bar{C}^g, \bar{C}^h \rangle \neq \bar{X}$ . We conclude that  $\hat{W}_i$  is normalized by  $N_A(V_i)$ . Therefore  $\hat{W}_i = [\bar{X}, N_A(V_i)]$ , and as  $\bar{W}_i = [\bar{X}, Y_0]$  we must have  $\bar{W}_i = \hat{W}_i$ . As  $Y_0$  acts irreducibly on  $\bar{W}_i/\bar{V}_i$  and  $\bar{V}_i < \bar{C} \cap \bar{C}^g \leq Z(\bar{W}_i)$  we must have  $\bar{W}_i$  Abelian as claimed.

Now we consider the case  $n = 3$ . First suppose that  $q \neq 4$ . Then  $\widehat{N_A(V_i)}$  contains a cyclic subgroup  $1 \neq Y_1$  of order  $q - 1$  or  $\frac{1}{3}(q - 1)$  with  $[L_i, Y_1] = 1$ . Then  $[\bar{X}, Y_1]$  is  $L_i$ -invariant and we conclude that  $[\bar{X}, Y_1] = \bar{W}_i^{L_i}$ . We have  $\bar{W}_i/\bar{V}_i$  isomorphic to  $\bar{V}_i$  as  $L_i$ -modules. Let  $\bar{Z}/\bar{V}_i$  be a 1-space of  $\bar{W}_i/\bar{V}_i$ .

Let  $Z_0, Z_1$  be as before, so that  $\bar{C}/\bar{Z}_1$  is abelian. If  $q = 4$  we also have  $\bar{C}/\bar{Z}_0$  abelian by [6, (2.2)]. So in this case  $\bar{C}$  is abelian. Suppose  $q > 4$  and let  $P = L_i \cap V$ . We may assume  $P$  centralizes  $\bar{C}/\bar{V}_i$ , so  $P$  centralizes  $\bar{W}_i/\bar{C}$ ,  $\bar{Z}_0$ , and  $\bar{V}_i/\bar{Z}_0$ . Note that  $[\bar{W}_i, P]$  covers  $\bar{C}/\bar{Z}_0$ . Now  $[\bar{W}_i, [\bar{W}_i, P], P] \leq [\bar{V}_i, P] \leq \bar{Z}_0$ , and  $[[\bar{W}_i, P], P, \bar{W}_i] \leq [\bar{V}_i, \bar{W}_i] = 1 \leq \bar{Z}_0$ . Consequently  $[[\bar{W}_i, P], [\bar{W}_i, P]] \leq \bar{Z}_0$  and again  $\bar{C}/\bar{Z}_0$  is abelian. Also  $P$  centralizing  $\bar{W}_i/\bar{C}$  and  $\bar{C}/\bar{V}_i$  implies  $P$  centralizes  $[\bar{C}, \bar{W}_i]$ . Thus  $[\bar{C}, \bar{W}_i] \leq \bar{Z}_0$ . Now let  $\bar{I}/\bar{V}_i$  be a 1-space of  $\bar{W}_i/\bar{V}_i$  complementing  $\bar{C}/\bar{V}_i$  and let  $\bar{I}_0 = [\bar{I}, t]$ . Then symmetry gives  $[\bar{I}, \bar{C}] \leq \bar{I}_0 \cap \bar{Z}_0 = 1$ . Since  $\bar{W}_i = \langle \bar{C}, \bar{I} \rangle$  we have the claim.

Finally we consider the case  $n = 3$  and  $q = 4$ . Let  $\bar{Y} = [\bar{X}, D] = \bar{Y}_1/\bar{V}_i \times \bar{Y}_2/\bar{V}_i$  with each factor  $D$ -invariant of order 4. As before  $\bar{Y}_1$  and  $\bar{Y}_2$  are abelian. But since  $Y_0$  is fixed-point-free on  $\bar{W}_i$ , an easy calculation in the Lie ring (or

use of commutators) shows  $\bar{Y}' < \bar{V}_i$ . If  $\bar{Y}' = 1$  we argue that  $\bar{Y}$  is characteristic in  $\bar{X}$ , forcing  $\bar{Y} = \bar{W}_i$ . Suppose  $\bar{Y}' \neq 1$ . Then regarding  $\bar{V}_i$  as an  $\mathbb{F}_2$ -space of dimension 4,  $\bar{Y}'$  is a 2-space in  $\bar{V}_i$ . Moreover we may choose  $g \in L_i$  with  $Y' \cap (Y')^g = 1$ . But consider  $Y \cap Y^g$ . This group is of index at most 2 in  $Y$ . But  $(Y \cap Y^g)' \leq Y' \cap (Y')^g = 1$ , so  $(Y \cap Y^g)' = 1$ . One checks that  $Y$  can have no such subgroup. This contradiction completes the proof of the claim.

Suppose  $g \in N(V_i \langle t \rangle) \leq N(V_i)$ . Then  $\bar{W}_i \cap \bar{W}_i^g$  has index of at most 2 in  $\bar{W}_i$  and is centralized by  $\langle \bar{W}_i, \bar{W}_i^g \rangle$ . As  $C_X(t) = V_i \langle t \rangle$  we must have  $\bar{W}_i = \bar{W}_i^g$ . So  $\bar{W}_i \trianglelefteq \bar{N}_G(\bar{V}_i)$ . Next write  $N(V_i \langle t \rangle) = O(N(V_i \langle t \rangle))(N(V_i \langle t \rangle) \cap N(X))$ . Since  $N(V_i \langle t \rangle) \cap N(X) \leq XC(t)$ , we have  $L_i \leq N(V_i \langle t \rangle) \cap N(X)$ , and we conclude that  $L_i \leq N(W_i)$ .

It remains to show that  $W_i$  is elementary Abelian. Suppose not. Then  $W_i$  is homocyclic of exponent 4 and we may take  $i = 1$ . Notice that if  $S_1 \in \text{Syl}_2(N_G(V_1)) \cap N_G(W_1 \langle t \rangle)$  then  $[S_1, t] = V_1$ . Let  $g \in N(S_1)$ . Then  $W_1^g < S_1$  and  $[W_1^g, t] \leq W_1^g \cap V_1$  is elementary Abelian. So  $t \in C_{S_1}(W_1^g/V_1) = W_1^g \langle t^g \rangle$ . Also  $t \notin S_1'$ , so  $t \notin V_1^g$ . Therefore  $t \in W_1^g t^g$  and so  $[S_1, t] \geq V_1^g$ . By order consideration  $V_1 = V_1^g$ . So  $g \in N(V_1)$ ,  $g \in N(C_S(V_1)) = N(W_1 \langle t \rangle)$ , and  $S_1 \in \text{Syl}_2(G)$ .

Now  $S_1$  normalizes  $V_2^k$ ,  $W_2^k$ , and  $(W_2 \langle t \rangle)^k$  for some  $k \in G$ . Suppose that  $W_2$  is elementary Abelian. Then  $[W_2^k, W_1] \leq W_2^k \cap W_1 \leq V_1$ , so  $W_2^k$  centralizes  $W_1/V_1$  forcing  $W_2^k \leq W_1 \langle t \rangle$ , which is impossible. Thus  $W_2$  is also homocyclic of exponent 4. As above  $[S_1, t] = V_1$  implies that  $t \in W_2^{k t^k}$  and  $V_2^k = V_1$ . Then  $W_2^k = W_1$ ,  $S_1$  and  $S_1^k \leq N(V_1)$ , and we may take  $k \in N(S_1)$ . This implies that  $W_2 = W_1^{k^{-1}} \trianglelefteq S_1$  and  $V_2 \trianglelefteq S_1$ . But then  $[V_2, W_1] \leq V_2 \cap W_1 \leq V_1$ , whereas  $V_2$  does not centralize  $W_1/V_1$ . This contradiction establishes (2.5).

(2.6) Let  $N_i = N_G(W_i)$  and  $C_i = V \cap L_i$ . Then  $S_i = W_i C_i \langle zw, t \rangle \notin \text{Syl}_2(N_i)$ .

*Proof.* Suppose that  $S_i \in \text{Syl}_2(N_i)$ . We may take  $i = 1$ . First assume that  $n = 3$ . We claim that  $W_1$  is weakly closed in  $S_1$ . For suppose  $W_1 \neq W_1^g \leq S_1$ . The elementary Abelian subgroups of maximal order in  $S_1/W_1$  have order  $2q$ , while  $|W_1^g| = q^4$ . Since  $q > 2$ ,  $|W_1^g \cap W_1| > q^2$ . Viewing  $W_1/V_1$  and  $V_1$  as  $\mathbb{F}_q$ -modules, this implies that either  $W_1^g \cap V_1$  generates  $V_1$  or  $(W_1^g \cap W_1) V_1/V_1$  generates  $W_1/V_1$ . But  $W_1^g$  centralizes  $W_1^g \cap W_1$ , so  $W_1^g \cap W_1 C_1 \langle t \rangle$  centralizes  $W_1/V_1$  or  $V_1$ , forcing  $W_1^g \cap W_1 C_1 \langle t \rangle \leq W_1 \langle t \rangle$ . Then  $W_1^g \leq C(W_1^g \cap W_1 C_1 \langle t \rangle)$  implies  $W_1^g \leq W_1 \langle t \rangle$ . As all involutions in  $W_1 \langle t \rangle$  are in  $W_1$  or  $V_1 t$ ,  $W_1^g = W_1$ , a contradiction. This proves the claim. In particular, since we are assuming  $S_1 \in \text{Syl}_2(N(W_1))$ , we must have  $S_1 \in \text{Syl}_2(G)$ . We can now obtain a contradiction by applying [4, Theorem 4]. Let  $q = 2^a > 2$ . The number  $r$  in Goldschmidt's result is at most  $a + 1$  as  $L_1 \cong SL(2, q)$ . Let  $x \in S_1 - W_1$  be an involution. Then  $[W_1, x] \geq q^2 = 2^{2a} > 2^r$ . So by [4, Theorem 4],  $W_1$  is strongly closed in  $S_1$ . The main theorem of [4] gives a contradiction.

We now assume that  $n \geq 4$ . First we construct a certain subgroup of  $G$ , and for this we introduce notation from the  $(B, N)$  structure of  $A$ . Let  $\mathcal{E}$  be a root system of type  $A_{n-1}$  with base  $\pi = \{\alpha_1, \dots, \alpha_{n-1}\}$ . We may assume that  $V$  is written  $V = \prod_{r \in \Sigma^+} U_r$  and generated by the root subgroups  $U_{\alpha_1}, \dots, U_{\alpha_{n-1}}$ . In this notation we may assume  $V_1 = U_{\alpha_1} U_{\alpha_1 + \alpha_2} U_{\alpha_1 + \dots + \alpha_{n-1}}$  and  $L_1 = \langle U_{\pm \alpha_2}, \dots, U_{\pm \alpha_{n-1}} \rangle$ . If  $n \geq 5$  set  $X = \langle U_{\pm \alpha_2} \rangle \cong SL(2, q)$  and if  $n = 4$  let  $X = A \cap C(\langle U_{\pm \alpha_2}, U_{\pm(\alpha_1 + \alpha_2)} \rangle)$ . In the latter case  $X$  is cyclic of order  $q - 1 > 1$ . Viewing  $L_1$  as matrices (via the modules  $V_1$  and  $W_1/V_1$ ) operating on the right we have

$$C_1 = \left\{ \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ * & & 1 \end{pmatrix} \right\},$$

$$U_{\alpha_2} = \left\{ \begin{pmatrix} 1 & & & \\ & \ddots & & \\ & & 1 & \\ & & & a1 \end{pmatrix} : a \in \mathbb{F}_q \right\}, \dots, U_{\alpha_{n-1}} = \left\{ \begin{pmatrix} 1 & & & \\ & a1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} : a \in \mathbb{F}_q \right\}.$$

In addition, if  $n \geq 5$  we set

$$J = \left\{ \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ * & \cdots & * & 1 & \\ * & \cdots & * & 0 & 1 \end{pmatrix} \right\} \quad \text{and} \quad F = \left\{ \begin{pmatrix} I & & \\ & 1 & \\ & & 1 \end{pmatrix} : I \in SL(n-3, q) \right\}.$$

Finally if  $n = 4$ , then  $X$  induces

$$\left\{ \begin{pmatrix} 1 & & \\ & 1 & \\ & & \alpha \end{pmatrix} : \alpha \in \mathbb{F}_q^\times \right\}$$

on  $V_1$  and we set

$$J = \left\{ \begin{pmatrix} 1 & & \\ & 1 & \\ * & * & 1 \end{pmatrix} \right\}, \quad F = \left\{ \begin{pmatrix} I & \\ & 1 \end{pmatrix} : I \in SL(2, q) \right\}.$$

Notice that  $F = \langle U_{\pm \alpha_1}, \dots, U_{\pm \alpha_{n-1}} \rangle$ . Now the construction of  $X$  forces  $X \leq C(U_{\alpha_1 + \dots + \alpha_{n-1}})$  so  $X$  centralizes an involution  $t \neq t^\theta \in tU_{\alpha_1 + \dots + \alpha_{n-1}}$ . We note that  $E(C(X) \cap C(t)) \cong SL(n-2, q)$  if  $n \geq 5$  and  $E(C(X) \cap C(t)) \cong SL(3, q)$  if  $n = 4$ . Moreover  $E(C(X) \cap C(t))$  is standard in  $C(X)$  and  $\langle t \rangle \in \text{Syl}_2(C(X) \cap C(E(C(X))))$ . Inductively either  $t \in Z^*(C(X))$ ,  $A \cong L_5(2)$  and



$E(C(X)) \cong J_2$ ,  $A \cong L_6(2)$  and  $E(C(X)) \cong A_{10}$ , or  $E(C(X))/Z(E(C(X))) \cong L(m, q^2)$  or  $L(m, q) \times L(m, q)$ , where  $E(C_A(X))/Z(E(C_A(X))) \cong L(m, q)$ . However, we know that  $X \leq C_A(t^q) = C_A(t^q)^{(\infty)} \leq N(A^q)^{(\infty)} = A^q$ , and consideration of  $C_{A^q}(X)$  shows that  $t \notin Z^*(C_{A^q}(X))$ . Consequently  $t \notin Z^*(C(X))$ .

If  $n = 4$  let  $X_1 = X$  and if  $n \geq 5$  let  $X_1 \leq X$  be a cyclic group of order  $q + 1$ . Write  $V_0 = C_{V_1}(X_1)$  and note that as  $X$  acts in the same way on  $W_1/V_1$  and on  $V_1$ ,  $|C_{W_1}(X_1)| = |V_0|^2$ . In case  $n \geq 5$  the above arguments show that  $E(C_G(X_1)) = E(C_G(X))$ , so  $W_0 = C_{W_1}(X_1) \leq E(C_G(X))$ . An easy argument shows  $W_0 \not\leq S_{10}$ . Suppose  $E(C(X)) \cong J_2$ . The Sylow 2-subgroups of  $J_2$  are isomorphic to the Sylow 2-subgroup of  $L(3, 4)\langle\sigma\rangle$  for  $\sigma$  a graph-field automorphism (see [21, (10.1)]. Let  $I \in \text{Syl}_2(E(C(X)))$  with  $t \in N(I)$  and  $W_0 \leq I$ . Now  $W_0$  is elementary of order  $2^4$  and  $W_0^t = W_0$ , so  $W_0 \leq I_0 \in \text{Syl}_2(L(3, 4))$ ,  $I_0 \trianglelefteq I\langle t \rangle$ . Looking in  $N_G(V_1) \cap C(X)$  we see that  $V_0$  is contained in  $E(C(X\langle t \rangle))$ , so from [21, (10.1), (4), (5)] we conclude that  $V_0 = Z(I_0)$ . Then  $I_0\langle t \rangle = I\langle t \rangle \cap C(V_0)$  and choosing  $h \in I - I_0$  we have  $[W_0, t] = V_0 = [W_0^h, t]$ . But then  $[t, I_0] = V_0$  and  $|E(C(X)) \cap C(t) \cap C(V_0)| \geq 2^4$ , against the structure of  $E(C(X)) \cap C(t) \cong \text{PGL}(2, 7)$ .

If  $n = 4$ ,  $V_0 = U_{\alpha_1+\alpha_2}U_{\alpha_1+\alpha_2+\alpha_3}$  and if  $n \geq 5$ ,  $V_0 = U_{\alpha_1+\alpha_2+\alpha_3} \cdots U_{\alpha_1+\cdots+\alpha_{n-1}}$ .  $V_0$  is the unique hyperplane, respectively  $((n-1)-2)$ -space, of  $V_1$  normalized by  $S_1$ . Then  $V_0 \leq E(C(X) \cap C(t))$  and by order  $O_2(C(X) \cap C(V_0)) = W_0$ . Moreover writing  $|V_0| = q^b$ ,  $N(W_0) \cap E(C(X))$  induces  $SL(b, q^2)$  or  $SL(b, q) \times SL(b, q)$  on  $W_0$ .

We claim that  $W_0 = C_{W_1}(X)$  and  $W_0 \leq S_1$ . By earlier remarks  $W_0 \leq C_{W_1}(X) \leq C_{W_1}(X_1) = W_0$ , so we have the first part of the claim.

For the second part of the claim we must show that  $W_0 \leq S_1$ . First assume that  $n \geq 5$ . Note that  $X \times F$  acts on  $W_1$  and on  $W_0 = C_{W_1}(X)$ . Also  $[X_1, W_1]$  is  $F$ -invariant and  $F$  centralizes  $[X_1, W_1]/[X_1, V_1]$  and also  $[X_1, V_1]$ . Therefore  $O^2(F)$  centralizes  $[X_1, W_1]$ . Except in the case  $A \cong L_5(2)$  this implies that  $F \leq C([X_1, W_1])$ . Assume also that  $A \not\cong L_5(2)$ . Then  $U_{\alpha_4} \leq F$  centralizes  $[X_1, W_1]$  and from the structure of  $E(C(X)) \cap N(W_0)$ ,  $||[U_{\alpha_4}, W_0]|| = q^2$ . So  $||[U_{\alpha_4}, W_1]|| = q^2$ , as  $W_1 = W_0 \times [W_1, X_1]$ . Now  $L_1$  is transitive on its root subgroups.  $J$  centralizes  $V_0$ ,  $V_1/V_0$ , and  $W_0V_1/V_1$ . If  $\alpha \in \Sigma^+$  with  $U_\alpha \leq J$ , then by the above  $[W_1, U_\alpha] \cap V_1 \leq V_0$ . In particular  $[W_0, U_\alpha] \leq V_0$  and  $U_\alpha \leq N(W_0)$ . As  $J$  is a product of such subgroups  $U_\alpha$ , we have  $J \leq N(W_0)$ . Hence  $C_1 \leq N(W_0)$ . As  $S_1 = W_1C_1\langle w, t \rangle$  and  $w$  normalizes  $X$ , we have  $W_0 \leq S_1$ , proving the claim in the cases  $n \geq 5$  and  $A \not\cong L_5(2)$ .

Suppose that  $A \cong L_5(2)$ . As above we only need show that  $||[U_{\alpha_4}, W_1]|| = q^2 = 4$ . Since  $U_{\alpha_2}$  and  $U_{\alpha_4}$  are  $L_1$ -conjugate it suffices to show that  $||[U_{\alpha_2}, W_1]|| = 4$ . Now  $D = \langle U_{\pm\alpha_2} \rangle \cong S_3$  and  $W_1 = W_0 \times [W_1, O_3(D)]$ . On the second factor  $\langle U_{\pm\alpha_2} \rangle = \bar{U}_{\alpha_2}O_3(D)$  acts with  $O_3(D)$  fixed-point-free. It follows that  $U_{\alpha_2}$  induces two copies of the regular representation on  $[W_1, O_3(D)]$ . As  $U_{\alpha_2} \leq C(W_0)$ ,  $||[U_{\alpha_2}, W_1]|| = 4$  as needed.

Finally we take the case  $n = 4$ . Write  $W_1 = W_0 \times [W_1, X]$  and note that

here  $[W_1, X]$  has order  $q^2$ . Also  $\langle U_{\pm\alpha_3} \rangle$  centralizes  $[W_1, X]/[V_1, X]$  and  $[V_1, X]$ , so  $\langle U_{\pm\alpha_3} \rangle = O^2(\langle U_{\pm\alpha_3} \rangle) \leq C([W_1, X])$ . Therefore  $|[W_1, U_{\alpha_3}]| = q^2$  and we argue as before. At this point the claim is proved.

Consider  $N_G(W_0) \geq S_1$ . Then  $C(W_0) \geq JX$  and  $[J, X] = J$ . As  $X \leq C(W_0)$ , it follows that  $J \leq C(W_0)$ . So  $S_1$  induces  $(S_1 \cap F)\langle w_1, t \rangle$  on  $W_0$ .

Assume that  $n \geq 5$ . Also assume that  $S_1 \cap C(W_0) \notin \text{Syl}_2(C(W_0))$ . Then there is an element  $g \in N(S_1) - S_1$  such that  $g \in C(W_0)$  and  $g^2 \in S_1$ . So  $V_0 = V_0^g$  and  $[t^g, S_1] = V_1^g \geq V_0$ . If  $V_1^g \leq W_1$ , then  $W_1 \leq C_{S_1}(V_1^g) = W_1^g$ , so  $g \in N(W_1)$ , against our hypothesis that  $S_1 \in \text{Syl}_2(N(W_1))$ . Therefore  $V_1^g \not\leq W_1$ . Now  $t^g \in S_1 \cap C(V_0) = \langle t \rangle W_1 J U_{\alpha_2}$ . Also  $t^g \notin W_1 t$  or else  $g \in W_1 C(t)$ . Say  $t^g$  induces an involution on  $W_1/V_1$  having  $l$  nontrivial Jordan blocks. Then  $[t^g, W_1/V_1] = q^l$ , and since  $V_0 \leq [t^g, S_1] \leq V_1^g \leq W_1$ ,  $l = 1$ . Also  $[t^g, S_1] \triangleleft S_1$ , so  $([t^g, S_1] \cap W_1) V_1/V_1 \leq S_1/V_1$ . It follows that  $[t^g, W_1] V_1/V_1$  is the unique 1-space of  $W_1/V_1$  fixed by  $S_1$ . Now  $W_1^g \leq S_1 \cap C(V_0) = W_1 J U_{\alpha_2}$ ,  $[W_1^g, t^g] = V_1^g$ , and  $W_1^g \leq S_1$ . It follows that  $W_1^g \leq W_1 J$  and so  $[W_1^g, W_1/V_1]$  is a hyperplane in  $W_1/V_1$ . Similarly  $[W_1^g, V_1]$  is a hyperplane in  $V_1$ . As  $[W_1^g, W_1] \leq W_1 \cap W_1^g$ ,  $|W_1 \cap W_1^g| \geq q^{2(n-2)}$ . Hence  $|W_1^g W_1/W_1| \leq q^2$ . But  $|[W_1^g, S_1] W_1/W_1| \geq |J|^{1/2} = q^{n-3} \geq q^2$ , and since  $W_1^g W_1/W_1 > [W_1^g, S_1] W_1/W_1$ , this is a contradiction. Therefore  $S_1 \cap C(W_0) \in \text{Syl}_2(C(W_0))$ .

However,  $S_1 \notin \text{Syl}_2(N(W_0))$ . In fact  $N(W_0) \cap C(X)$  contains a  $\langle t, w \rangle$ -invariant subgroup  $H$  with  $H$  inducing  $SL(n-3, q^2)$  or  $SL(n-3, q) \times SL(n-3, q)$  on  $W_0$ . Then  $S_1 \in C(W_0) H \langle t, w \rangle$  and there is an element  $g \in N(S_1) \cap N(W_0)$  with  $g \notin S_1$ ,  $g^2 \in S_1$ , and  $t^g \in t U_{\alpha_3}^\#(S_1 \cap C(W_0))$ . As before, we must have  $V_1^g \leq W_1$  and  $([S_1, t^g] \cap W_1) V_1/V_1 \leq S_1/V_1$ . It follows that  $[S_1, t^g] V_1/V_1 \cap W_1/V_1$  contains an  $(n-3)$ -subspace of  $W_1/V_1$ . Similarly  $[S_1, t^g] \cap V_1$  contains an  $(n-3)$ -subspace of  $V_1$ . So  $|V_1^g \cap W_1| \geq q^{2(n-3)} \geq q^{n-1} = |V_1| = |V_1^g|$ . As  $V_1^g \leq W_1$ , this is a contradiction.

Finally consider the case  $n = 4$ . Choose  $g \in N(S_1) \cap N(W_0)$  with  $g^2 \in S_1$ , but  $g \notin S_1$ . First assume that  $S_1 \cap C(W_0) \notin \text{Syl}_2(C(W_0))$  so that  $g$  may be chosen in  $C(W_0)$ . Then  $V_0^g = V_0$  and  $t^g$  does not centralize  $W_1/V_1$ . So  $[t^g, W_1/V_1] = q$  and  $|V_1^g \cap W_1| \geq |V_0| q = q^3 = |V_1^g|$ . But as before  $V_1^g \leq W_1$ , so this is a contradiction. Therefore  $S_0 = S_1 \cap C(W_0) \in \text{Syl}_2(C(W_0))$ . We consider the action of  $N(S_0) \cap N(W_0)$  on  $S_0/W_0$ . We know that  $|C_{S_0/W_0}(t)| = q^3$  and  $|S_0/W_0| = q^4$ . Also  $N(W_0) \cap N(S_0)$  contains a subgroup  $(B_1 \times B_2)\langle t \rangle$  with  $B_1 \cong B_2 \cong SL(2, q)$  and  $B_1^t = B_2$ . So any  $b_1, b'_1$  in  $B_1$ ,  $\langle t, t^{b_1}, t^{b'_1} \rangle$  centralizes a subgroup of order at least  $q$  in  $S_0/W_0$ . But with proper choice of  $b_1, b'_1$  we have  $\langle t, t^{b_1}, t^{b'_1} \rangle = (B_1 \times B_2)\langle t \rangle$ . As we easily check that  $C_{S_0/W_0}(\langle U_{\pm\alpha_3} \rangle) = 1$ , this is impossible. The proof of (2.6) is now complete.

(2.7) Let  $N_i = N_G(W_i)$ .

(i)  $O(N_i^{(\infty)}) = Z(N_i^{(\infty)})$ .

(ii) Letting bars denote images in  $N_i/W_i$ , we have  $E(\bar{N}_i)/Z(E(\bar{N}_i)) \cong L_{n-1}(q^2)$  or  $L_{n-1}(q) \times L_{n-1}(q)$ .

*Proof.* Consider the group  $N'_i$ . We may take  $i = 1$ . Let  $X = O(N'_1)$ , let  $L_1 = \langle U_{\pm\alpha_1}, \dots, U_{\pm\alpha_{n-2}} \rangle$  and let  $z$  be an involution in  $U_{\alpha_1 + \dots + \alpha_{n-2}}$ . Then by (2.1)  $t \sim tz$ , so write  $tz = t^\theta$ . Also write  $X = C_X(t) C_X(tz) C_X(z)$ . As  $C_X(t) \leq C(W_1) \leq C(V_1)$ ,  $C_X(t) \leq O(N(A))$  and hence  $C_X(t) \leq C(A) \leq C(z)$ . Next consider  $C_X(tz) = C_X(t^\theta) \leq N(A^\theta)$ . By (2.1) the Sylow 2-subgroups of  $C_A(t^\theta)$  are contained in  $A^\theta$ . In particular  $C_{V_1}(t^\theta) = C_{V_1}(z)$  is a hyperplane in  $V_1$  and centralized by  $C_X(t^\theta)$ . Since  $C_L(t^\theta)$  normalizes  $C_X(t^\theta)$  we must have  $C_X(t^\theta) \leq O(N(A^\theta))$  and again  $C_X(t^\theta) \leq C(z)$ . Therefore  $X$  is centralized by  $z$ , so  $C_{N_1}(X)$ , being normal in  $N_1$ , must cover  $E(\langle z^{N_1} \rangle W_1 X / W_1 X)$ .

We know that  $W_1$  controls fusion of  $t^G \cap W_1 t$ . In fact the only involutions in  $W_1 \langle t \rangle - W_1$  are those in  $V_1 t = t^{W_1}$ . Therefore  $C_{N_1}(t) = \bar{L}_1 \langle \bar{w}, \bar{t} \rangle$  and it is easy to see that  $\bar{L}_i$  is standard in  $\bar{N}_i$ . Also  $\bar{L}_i \not\leq \bar{N}_i$  by (2.6).

By the main theorem in [6] if  $n = 3$ , Solomon's theorem [17] if  $n = 5$  and  $q = 2$ , or induction for all other values of  $n$  and  $q$  we conclude that either (2.7) holds, or  $n = 3$  and  $E(\bar{N}_1) \cong U_3(q)$ ,  $L_3(q)$ ,  $G_2(3)$ ,  $M_{12}$ , or  $n = 5$ ,  $q = 2$ , and  $E(\bar{N}_1) \cong A_{10}$ , or  $HS$ .

If  $E(\bar{N}_1) \cong M_{12}$ , then  $q = 4$  and  $|W_1| = q^4 = 2^8$ . But  $11 \nmid |M_{12}|$  implies that  $E(\bar{N}_1)$  must act trivially on  $W_1$ , impossible. If  $E(\bar{N}_1) \cong U_3(q)$ , then by [3, (4D)] we get the same contradiction. Suppose  $E(\bar{N}_1) \cong L_3(q)$ . Here  $\bar{t}$  acts as a graph automorphism, so  $E(\bar{N}_1) \langle \bar{t} \rangle$  cannot act faithfully on a 2-group of order less than  $q^6$ . Again we have a contradiction. If  $E(\bar{N}_1) \cong G_2(3)$ , then  $A \cong L_3(8)$  and  $N_A(V_1) \geq L_2(8) \times Z_7$ . Viewing  $N_A(V_1) \leq N_A(W_1)$  the  $Z_7$  factor must centralize  $E(\bar{N}_1)$  and this leads to  $C_2(3) \leq L_4(8)$ . Impossible.

Suppose  $n = 5$ ,  $q = 2$ . If  $E(\bar{N}_1) \cong A_{10}$ , let  $Y \leq A$  be a subgroup of order 5. Then  $C = C(Y) \cap E(\bar{N}_1)$  involves  $S_5$ . But this forces  $C$  to be irreducible on  $W_1$  and by Schur's lemma  $C$  is isomorphic to a subgroup of  $GL(2, 2^4)$ . This is impossible. Finally if  $E(\bar{N}_1) \cong HS$ , then  $11 \nmid |E(\bar{N}_1)|$ . As  $|W_1| = 2^8$ , this forces  $E(\bar{N}_1)$  to act trivially on  $|W_1|$ , a final contradiction.

### 3. $\widetilde{E(\bar{N}_i)} \cong L_{n-1}(q^2)$ FOR $i = 1$ OR $2$ .

In this section we make the assumption that  $\widetilde{E(\bar{N}_i)} \cong L_{n-1}(q^2)$  for  $i = 1$  or  $2$ , recalling from (2.7) that  $\bar{N}_i = N_i / W_i$ . We may assume that  $i = 1$ .

(3.1)  $E(\bar{N}_1) \cong SL(n-1, q^2)$  and  $W_1$  can be viewed as an  $\mathbb{F}_{q^2}(E(\bar{N}_1))$ -module on which  $E(\bar{N}_1)$  induces the usual representation of  $SL(n-1, q^2)$ . Also  $N_1^{(\infty)}$  splits over  $W_1$ . In particular the Sylow 2-subgroups of  $N_1^{(\infty)}$  are isomorphic to those of  $L(n, q^2)$ .

*Proof.* We begin by proving the first statement, and for this start with the case  $n \geq 4$ . We know that  $E(\bar{N}_1)$  is an image of  $SL(n-1, q^2)(23, \S 8)$  so consider  $W_1$  as an  $\mathbb{F}_2$ -module for  $SL(n-1, q^2)$ . Let  $q^2 = 2^a$ . By Steinberg [19],  $\mathbb{F}_{q^2}$  is

a splitting field for  $SL(n-1, q^2)$  so we have  $W_1 \otimes \mathbb{F}_{q^2} = M_1 \oplus \cdots \oplus M_r$  a sum of inequivalent algebraically conjugate, absolutely irreducible representations of  $SL(n-1, q^2)$ . If  $\theta: x \rightarrow x^2$  is the Frobenius map, then  $M_1$  and  $M_1^{\theta^r}$  are equivalent and we write  $a = rb$ . By the tensor product theorem [19],  $M_1 \cong T_1 \otimes T_2^{\theta} \otimes \cdots \otimes T_a^{\theta^{a-1}}$ , where the  $T_i$  are absolutely irreducible and we may assume  $T_1$  is nontrivial. As  $M_1 \cong M_1^{\theta^r}$ ,  $\dim(M_1) \geq (\dim(T_1))^b$ . Also since the Sylow 2-subgroups of  $SL(n-1, q^2)$  has class  $n-2$ ,  $\dim(T_1) \geq n-1$ . We now have  $a(n-1) = \dim_{\mathbb{F}_q}(W_1) = \dim_{\mathbb{F}_{q^2}}(W_1 \otimes \mathbb{F}_{q^2}) \geq r(\dim(T_1))^b \geq r(n-1)^b$ . So  $b(n-1) \geq (n-1)^b$ , and as  $n-1 \geq 3$  we conclude that  $b = 1$  and  $W_1 \otimes \mathbb{F}_{q^2} = M_1 \oplus M_1^{\theta} \oplus \cdots \oplus M_1^{\theta^{a-1}}$ , where  $M_1$  is the usual module for  $SL(n-1, q^2)$ . If  $V$  is the usual module for  $SL(n-1, q^2)$ , viewed as an  $\mathbb{F}_2$ -module, then  $V \otimes \mathbb{F}_{q^2}$  is also isomorphic to  $M_1 \oplus \cdots \oplus M_1^{\theta^{a-1}}$ . It follows that  $V \cong W_1$  as  $\mathbb{F}_2(SL(n-1, q^2))$  modules and the result holds.

Now suppose  $n = 3$ . The above arguments carry over and the relevant inequality is  $b2 \geq 2^b$ . If  $b = 1$  there is no problem. So assume that  $b = 2$ . Then in the earlier notation  $M_1 \cong V \otimes V^q$  where  $V$  is the usual module for  $SL(2, q)$ . We must show that this does not occur. Let  $S_0 \in \text{Syl}_2(N_1^{(\infty)})$  and let  $H_1$  be a  $t$ -invariant 2-complement in  $N_{N_1^{(\infty)}}(S_0)$ . So  $H_1$  is cyclic of order  $q^2 - 1$ . The action of  $H_1$  on  $M_1$  shows that  $\bar{H}_1^{q+1}$  has nontrivial centralizers on  $M_1$ . Hence  $C_{W_1}(\bar{H}_1^{q+1}) \neq 1$ . However  $\bar{H}_1^{q+1}$  is covered by  $\overline{C_{\bar{H}_1}(t)}$  and  $C_{\bar{H}_1}(t)$  acts in the same way on  $V_1$  and on  $W_1/V_1$ . As  $C_{\bar{H}_1}(t)$  is fixed-point-free on  $V_1$ , this is impossible.

To complete the proof of (3.1) it suffices by Gaschutz' result [9, 17.4] to show that the Sylow 2-subgroups of  $N_1^{(\infty)}$  split over  $W_1$ . So consider the group  $N_1^{(\infty)}$  and note that  $O(N_1^{(\infty)}) = 1$ . Let  $S_0 \in \text{Syl}_2(N_1^{(\infty)})$  be such that  $V \leq S_0$  and  $S_0$  is  $\langle t \rangle$ -invariant. Viewing  $W_1$  as an  $\mathbb{F}_{q^2}$ -module let  $W_0$  be the 1-space of  $W_1$  stabilized by  $S_0$  and let  $L = C_{S_0}(W_1/W_0)$ . Then  $N(L) \cap N_1^{(\infty)} = LDH$  where  $\bar{D} \cong SL(n-2, q^2)$ ,  $[\bar{D}, \bar{H}] = 1$ ,  $Z(\bar{D}) \leq \bar{H}$  is cyclic of order  $q^2 - 1$ , and  $C_L(\bar{H}) = 1$  or  $W_0$ , the latter occurring only if  $\bar{H} = Z(\bar{D})$ . We may take  $H$  to be cyclic of order  $q^2 - 1$ , so the Frattini argument implies that  $N(L) \cap N_1^{(\infty)} \leq LN(H)$  and  $L \cap N(H) = 1$  or  $W_0$ . Checking multipliers we see that  $(N(H) \cap N_1^{(\infty)})$  complements  $L$  in  $LD$  unless  $\bar{D} \cong SL(3, 4)$ . So except for this one case, which we handle later, we may assume  $DH \cong \bar{D}\bar{H}$ .

Recall the subgroups  $V_1 \leq W_1$  and  $V_2$  from Section 2. We have  $V_2 \leq W_1L$  and  $V_2 \cap W_1 = V_1 \cap V_2 \leq W_0$ . Observe that  $L/W_0$  is elementary Abelian. Indeed  $L/W_0$  contains  $W_1/W_0$  as a normal subgroup and  $DH$  is transitive on  $L/W_1$ . Also  $W_1/W_0$  is central in  $L/W_0$  and  $L - W_1$  contains involutions, as  $V_2 \not\leq W_1$ . Therefore each coset of  $W_1/W_0$  in  $L/W_0$  contains involutions, and  $L/W_0$  is elementary Abelian. Suppose  $n = 3$ . Then  $q \geq 4$  and there is a subgroup  $H_1 \leq N_1^{(\infty)}$  such that  $H_1 \leq N(L)$  and  $\bar{H}_1$  is cyclic of order  $q^2 - 1$ . Also  $L = S_0$ . Now  $H_1$  acts on  $L/W_0$ , and  $L/W_0$  is the sum of inequivalent  $\mathbb{F}_2(H_1)$ -submodules,  $L/W_0 = W_1/W_0 \times C/W_0$ . One checks that  $C/W_0$  and  $W_0$  are inequivalent as  $\mathbb{F}_2(H_1)$ -modules, so  $C$  is not homocyclic. As  $|C/W_0|$  is a square,

$C$  is not a Suzuki 2-group [3] and it follows that  $C$  is elementary Abelian. Then  $S_0$  splits over  $W_1$  as required. We now assume  $n \geq 4$ .

Assume  $q > 2$ . Then in  $N_A(V_1)$  there is a nontrivial subgroup  $H_1$  with  $|H_1| \mid q-1$  and centralizing  $L_1 \cong SL(n-1, q)$ . We may choose  $H_1 \leq N(W_1) = N_1$  and it follows that  $\bar{H}_1 \leq C_{N_1}(\bar{N}_1^{(\infty)})$ . As  $H_1$  is fixed-point-free on  $V_1$ , it is also fixed-point-free on  $W_1$ . An application of the Frattini argument now implies that  $N_1 = W_1 N_{N_1}(O(N_1) H_1)$ , a semidirect product. Again we have the result.

Finally assume  $q = 2$ . Then  $n \geq 5$ ,  $|V_2| = 2^{n-1}$ , and  $|V_2/V_2 \cap W_1| = 2^{n-2}$ . For each involution  $x \in V_2 - W_1$ ,  $[L, x] = W_0$  and so  $|C_L(x)| = \frac{1}{4}|L|$ . Therefore  $|C_L(V_2)| \geq (\frac{1}{4})^{n-2}|L| = (\frac{1}{4})^{n-2}4^{n-3} = 4^{n-1} = |W_2|$ . Also  $C_L(V_2) \cap W_1 = W_0$ , so  $L = W_1 C_L(V_2)$ . The group  $L_1 \cap L_2 \cong SL(n-2, q)$  normalizes  $C_L(V_2)$  inducing contragredient representations on  $V_1/V_1 \cap V_2$  and  $V_2/V_1 \cap V_2$ . As  $W_2 \langle t \rangle \in \text{Syl}_2(C(V_2))$  and  $|C_L(V_2)| = |W_2|$ , we conclude that  $L_1 \cap L_2$  acts on  $C_L(V_2)/W_0$  as the sum of two representations equivalent to  $V_2/V_1 \cap V_2$ . Now let  $J \leq L_1 \cap L_2$  be a Singer cycle. So  $J$  is cyclic of order  $2^{n-2} - 1$ . From the above facts it follows that  $L/W_0$  decomposes as the sum of four irreducible  $\mathbb{F}_2(J)$ -submodules;  $W_1/W_0$  is the sum of two equivalent  $\mathbb{F}_2(J)$ -submodules and  $L/W_0 = W_1/W_0 \times X/W_0$ , where  $X/W_0 \geq V_2 W_0/W_0$  and is the sum of two irreducible submodules contragredient (thus inequivalent) to those in  $W_1/W_0$ . Now  $\bar{J} \leq \overline{L_1 \cap L_2} \leq \bar{D}$  so  $\bar{J}$  is centralized by a Singer cycle in  $\bar{D}$ , having order  $4^{n-2} - 1$  or  $\frac{1}{3}(4^{n-2} - 1)$ . This group then normalizes  $X$  as does  $L_1 \cap L_2$ . Therefore  $D \leq N(X)$  (see [14]) and since  $D$  is transitive on  $X/W_0$ ,  $X$  must be elementary Abelian. As  $|X| = |W_2|$ ,  $X = W_2$ . We now conclude that  $E(N_2/W_2 O(N_2)) \cong SL(n-1, q^2)$  and the action on  $W_2$  is as on the usual module. This determines the action of  $D$  on  $X$  and so  $X = W_0 \times X_1$ , where  $X_1$  is  $D$ -invariant. So a Sylow 2-subgroup of  $X_1 D$  complements  $W_1$  in a Sylow 2-subgroup of  $N_1^{(\infty)}$ , except in the one possible case of  $\bar{D} \cong SL(3, 4)$ .

In the exceptional case  $\bar{H} \leq \bar{D}$ . Suppose  $N_1$  contains a subgroup inducing  $GL(4, 4)$  on  $W_1$ . Then we easily see that  $N_1^{(\infty)}$  splits over  $W_1$ , as desired. So we suppose this is not the case.

Let  $S \in \text{Syl}_2(W_1 W_2 D)$  with  $V \leq S$ . Choose  $I \leq N_1^{(\infty)}$  elementary of order  $3^3$  and such that  $\bar{I} \leq N(\bar{S})$  is the Cartan subgroup with respect to the root system  $\Sigma_1$  spanned by  $\{\pm\alpha_2, \pm\alpha_3, \pm\alpha_4\}$ . Let  $r \in \Sigma_1$  and  $Z_r$  the root subgroup of  $N_1^{(\infty)}$  containing  $\bar{V}_r$ . Let  $I_0 = C_I(\bar{Z}_r)$ . One checks that  $I_0 \cong Z_3 \times Z_3$  and  $C_{W_1}(I_0) = 1$ . So if  $D_r$  is the preimage of  $Z_r$  we have  $D_r = W_1 Z_r$  and  $W_1 \cap Z_r = 1$ , where  $Z_r = C_{D_r}(I_0)$ . Each  $Z_r$  is  $I$ -invariant. Also viewing  $W_1$  as an  $\mathbb{F}_4$ -space we see that there are precisely four 1-spaces of  $W_1$  invariant under  $I$ . Label these  $Z_1 = Z_{\alpha_1}$ ,  $Z_2 = Z_{\alpha_1+\alpha_2}$ ,  $Z_3 = Z_{\alpha_1+\alpha_2+\alpha_3}$ , and  $Z_4 = Z_{\alpha_1+\alpha_2+\alpha_3+\alpha_4}$  in such a way that  $1 < Z_4 < Z_4 \cdot Z_3 < Z_4 \cdot Z_3 \cdot Z_2 < W_1$  is an  $IS$ -composition series of  $W_1$ . We then have  $S = \prod_{r \in \Sigma^+} Z_r$  and one checks that for  $r \neq s$  in  $\Sigma^+$ ,  $C_I(Z_r) \neq C_I(Z_s)$ . Now the proof of Lemma 3 in [22] shows that every  $I$ -invariant subgroup of  $S$  is a product of some of the root subgroups  $Z_r$ ,  $r \in \Sigma$ .

The earlier arguments show that  $I \leq N(W_2)$ . It follows that  $N_2^{(\infty)}/W_2 \cong SL(4, 4)$  and as  $[S, I] = S$ ,  $S \in \text{Syl}_2(N_2^{(\infty)})$ . If  $N_2$  contains a subgroup inducing  $GL(4, 4)$  on  $W_2$ , then as above  $N_2^{(\infty)}$  splits over  $W_2$ . It follows that  $W_0 D$  splits over  $W_0$ . So assume this is not the case. Then  $I \leq N_2^{(\infty)}$  and we carry through the above process of labeling root subgroups with respect to  $N_2^{(\infty)}$ . This labeling necessarily coincides with the earlier labeling since in each case the root subgroups in  $S$  are the minimal  $I$ -invariant subgroups of  $S$ .

Considering matrices in  $N_1^{(\infty)}$  we see that  $Z_{\alpha_2} \leq N(Z_{\alpha_1} Z_{\alpha_1 + \alpha_2})$ . So  $Y = \langle Z_{\alpha_1}, Z_{\alpha_2} \rangle = Z_{\alpha_1} Z_{\alpha_2} Z_{\alpha_1 + \alpha_2}$  and  $Y \cap W_2 = 1$ . It follows that  $W_2 Y$  is  $N_2$ -conjugate to  $W_2(S \cap D)$ , so by Gaschutz' theorem  $D$  splits over  $W_2$  and hence over  $L$  as needed.

$$(3.2) \quad (i) \quad E(\bar{N}_1) \langle \bar{w}, \bar{t} \rangle = E(\bar{N}_1) \langle \bar{w} \rangle \text{ and } E(\bar{N}_1) \cap \langle \bar{w} \rangle = 1.$$

(ii)  $t$  induces a field automorphism on  $E(\bar{N}_1)$ .

(iii) a Sylow 2-subgroup  $S_1$  of  $N_1$  has the form  $S_1 = S_0 \langle w_1 \rangle$ , where  $S_0 \in \text{Syl}_2(N_1^{(\infty)})$  and  $S_0 \cap \langle w_1 \rangle = 1$ . Also  $t \in \langle w_1 \rangle$  and we may take  $V \leq S_0$ .

*Proof.* (ii) follows from the structure of  $\text{Aut}(L(n-1, q^2))$ . By Lang's theorem [10] each involution in  $E(\bar{N}_1) \langle \bar{t} \rangle$  is  $E(\bar{N}_1)$ -conjugate to  $\bar{t}$ . It follows from this and  $t^g \cap tW_1 = tW_1$ , that  $N_1 = N_1^{(\infty)} C(t)$ . Considering  $\text{Aut}(L(n-1, q^2))$  we have (i). For (iii) first obtain an  $N_1$ -conjugate  $w_1$  of  $w$  with  $t \in \langle w_1 \rangle$ . Then choose  $S_1 \in \text{Syl}_2(N_1)$  with  $V \langle w_1 \rangle \leq S_1$  and set  $S_0 = S_1 \cap N_1^\infty$ .

(3.3) Let  $S_1 = S_0 \langle w_1 \rangle$ , where  $V \leq S_0 \in \text{Syl}_2(N_1^{(\infty)})$  and  $S_1 \in \text{Syl}_2(N_1)$ . There is an involution  $d \in N(S_1) - S_1$  such that  $S = S_1 \langle d \rangle \in \text{Syl}_2(G)$  and  $d \in C(t)$ .

*Proof.* First we show that  $S_1 \notin \text{Syl}_2(G)$ . For suppose otherwise. If  $t^G \cap S_0 = \emptyset$ , then transfer applies and we have  $t \notin G'$ . Then  $S_1 \cap G' = S_0$  (as  $S_1 \leq N_1^{(\infty)}$ ), so  $G'$  has Sylow 2-subgroups of type  $L_n(q^2)$ . By McBride [12],  $E(G'/O(G')) \cong L_n(q^2)$ . So by (1.3),  $E(G')/Z(G') \cong L_n(q^2)$ , whereas  $G$  is assumed to be a counterexample to the main theorem. Therefore  $t^G \cap S_0 \neq \emptyset$ . By (2.1),  $t^G \cap U = \emptyset$  and since  $W_1^\#$  is fused in  $N_1$ ,  $t^G \cap W_1 = \emptyset$ . Say  $t^g \in S_0 - W_1$  and  $t^g$  has precisely  $l$  nontrivial Jordan blocks on  $W_1$ , when  $W_1$  is viewed as an  $\mathbb{F}_{q^2}$ -module. Write  $S_0 = W_1 S_{00}$  where  $W_1 \cap S_{00} = 1$ . We may assume that  $S_0$  normalizes the  $l$ -space  $[W_1, t^g]$  and that a basis has been chosen for  $W_1$  in such a way that  $S_{00}$  is represented on  $W_1$  by lower triangular matrices and  $t^g$  is represented by the matrix

$$\begin{bmatrix} 1 & & & & & & \\ & \cdot & & & & & \\ & & \cdot & & & & \\ & & & \cdot & & & \\ 1 & & & & 1 & & \\ & \cdot & & & & \cdot & \\ & & \cdot & & & & \cdot \\ & & & \cdot & & & \\ & & & & 1 & & \\ & & & & & & 1 \end{bmatrix}.$$

Let  $t^g = sw$  with  $s$  an involution in  $S_0$  and  $w \in W_1$ . Then  $w \in C_{W_1}(t^g) = C_{W_1}(s)$ , which has dimension  $n - 1 - l$ . Set  $J = \langle s^{S_0} \rangle$ . Then  $C_{W_1}(s) = C_{W_1}(J)$ . Moreover  $C_{W_1}(t^g)$  and  $C_{W_1}(t^g)J$  are normal in  $S_0$ , so  $[S_0, t^g] \leq C_{W_1}(t^g)[J, S_0]$ .

One checks that  $[J, S_0] \leq C_{W_1}(t^g)[J, S_0]$  and  $|[J, S_0]| = (q^2)^{(1/2)l(l-1)} = q^{l(l-1)}$ . Therefore  $|[S_0, t^g]| \leq q^{l(l-1)}(q^2)^{n-l-1}$  and  $|C_{S_0}(t^g)| \geq q^a$  for  $a \geq n(n-1) - (l(l-1) + 2(n-l-1))$ . If  $n \geq 4$ , then  $|C_{S_0}(t^g)|$  has order strictly greater than the order of a Sylow 2-subgroup of  $C(t)$ . This is a contradiction. If  $n = 3$ ,  $C_{S_0}(t^g)$  contains an elementary subgroup of order  $q^4$ . Again this is impossible. So we must have  $S_1 \notin \text{Syl}_2(G)$  as claimed.

Choose  $d \in N(S_1) - S_1$  with  $d^2 \in S_1$ . As above we have  $t^G \cap S_0 = \emptyset$ , so  $t^d \in tS_0$ . As  $t$  induces a field automorphism on  $S_0$ ,  $t^d = t^s$  for  $s \in S_0$ , and we may take  $d \in C(t)$ . Then  $V\langle w, d \rangle \in \text{Syl}_2(N(A))$ . The structure of  $\text{Aut}(S_0)$  is known [11] and from the description there it is clear that  $t \notin (S_0\langle w, d \rangle)'$ . So either we may take  $d$  as an involution or  $w = t = d^2$ . Suppose the latter occurs. If  $S_0\langle d \rangle \in \text{Syl}_2(G)$ , then by the above argument  $t^G \cap S_0 = \emptyset$ . But then we use transfer and [12] to get a contradiction. If  $S_0\langle d \rangle \notin \text{Syl}_2(G)$ , then as above there exists a 2-element  $g \in C(t) \cap N(S_0\langle d \rangle)$  and  $g \notin S_0\langle d \rangle$ . This is impossible as  $V\langle w, d \rangle = V\langle d \rangle \in \text{Syl}_2(N(A))$ . We may now assume  $d^2 = 1$ .

Finally we must show that  $S_1\langle d \rangle \in \text{Syl}_2(G)$ . So suppose not and let  $s \in N(S_1\langle d \rangle) - S_1\langle d \rangle$ . We have seen that  $t^G \cap S_0 = \emptyset$ , so  $t^s \in tS_0, dS_0$ , or  $dtS_0$ . Each factor in the lower central series for  $S_0$  is a free  $\mathbb{F}_2\langle t \rangle$ -module, so it follows from (1.2) that each involution in  $S_0t$  is conjugate in  $S_0$  to  $t$ . As  $S_1$  contains  $V\langle w, d \rangle \in \text{Syl}_2(C(t))$  we conclude  $t^s \notin tS_0$ . Replacing  $d$  by  $dt$ , if necessary, we may assume  $t^s \in dS_0$ . Also we see that  $s$  normalizes  $S_0$  as follows. Note that  $Z = Z(S_0) = C(S_1\langle d \rangle)' \cap S_1\langle d \rangle$ , so  $s \in N(Z(S_0))$ . Next  $Z_2(S_0) = Z_2$  where  $Z_2/Z_1 = C((S_1\langle d \rangle)'Z_1/Z_1)$ . Continuing we get  $S_0 = Z_{n-1}$  to be  $s$ -invariant. It follows that each involution in  $dS_0$  is  $S_0$ -conjugate to  $d$  and we may assume that  $t^s = d$ .

Now  $d$  interchanges  $V_1, V_2$  and  $s \in N(S_0)$ . Choosing  $W_1, W_2 \leq S_0$  we have  $d$  interchanging  $W_1$  and  $W_2$ . From the structure of  $\text{Aut}(S_0)$  [11] we have a contradiction. So  $S_1\langle d \rangle \in \text{Syl}_2(G)$  and the proof of (3.3) is complete.

(3.4)  $G$  contains a normal subgroup  $G_0$  such that  $S_0\langle d_1 \rangle \in \text{Syl}_2(G_0)$ , where  $d_1 = d$  or  $dt$ .

*Proof.* Consider the transfer into  $S_1\langle d \rangle/S_0\langle d \rangle$ . To transfer out  $\langle w \rangle$  it suffices to show that  $t^G \cap S_0\langle d \rangle = \emptyset$ . Once this is accomplished, we have the result by using the fact that  $S_0 \leq N(S_0)' \leq G'$ .

Suppose then that  $t^G \cap S_0\langle d \rangle \neq \emptyset$ . As in (3.3)  $t^G \cap S_0 = \emptyset$ , so suppose  $t^g \in S_0d$ . Consider  $t^g$  acting on  $X = (N_1 \cap N_2)'$  and assume  $n \geq 5$ . Then  $X$  induces  $SL(n-2, q^2)$  on  $W_1W_2/W_1 \cap W_2$ . Replacing  $d$  by  $dt$ , if necessary, we may assume that  $d$  induces a graph-field automorphism on  $X/O_2(X)$ , so by [2, (19.6)] we have  $t$  conjugate to an involution in  $dW_1W_2$ . So take  $t^g \in dW_1W_2$ ,

and since  $W_1W_2/W_1 \cap W_2$  is a free  $\mathbb{F}_2\langle d \rangle$ -module, we may take  $t^g \in d(W_1 \cap W_2)$ . Since  $X \leq C(W_1 \cap W_2)$  and  $W_1W_2/(W_1 \cap W_2)$  is a free  $\mathbb{F}_2\langle d \rangle$ -module we have  $C(t^g)$  covering  $O^2(C(d) \cap (X/W_1W_2))$ . But this implies that  $E(C(t)) \sim E(C(t^g))$  contains a section isomorphic to  $O^2(SU(n-2, q))$ , which is impossible.

So  $n = 3$  or  $4$ . First assume that  $n = 4$ . Then  $X = (N_1 \cap N_2)'$  induces  $SL(2, q^2)$  on  $W_1W_2/W_1 \cap W_2$  and  $d$  or  $dt$  must centralize  $X/O_2(X)$ . Replacing  $d$  by  $dt$ , if necessary, we may assume  $t^g \in dS_0$  and  $[t^g, S_0], [d, S_0] \leq W_1W_2$ . As  $t^g$  interchanges  $W_1$  and  $W_2$ , the stabilizer in  $X$  of  $t^g(W_1 \cap W_2)$  covers  $X/O_2(X)$  and has order  $q^4 \mid |X/O_2(X)|$ . From the action of  $X$  on  $O_2(X)$  we conclude that  $C_X(t^g)$  covers  $X/W_1 \cap W_2$ . But  $C_G(t)$  contains no subgroup isomorphic to  $C_X(t^g)$ . This is impossible. Therefore  $n = 3$ . Choose a subgroup  $H \leq A$  with  $H$   $d$ -invariant,  $H$  cyclic of order  $q-1$ ,  $H$  fixed-point-free on  $V_1 \cap V_2$ . Then  $H$  is fixed-point-free on  $W_1 \cap W_2$ . Also  $[H, d] = 1$  or  $[H, d]$  has order  $q^{1/2} + 1$ , depending on whether  $C_A(d) \cong SL(2, q)$  or  $U_3(q^{1/2})$ . Considering  $H\langle d \rangle$  acting on  $W_1 \cap W_2$  we conclude that  $C(d) \cap W_1 \cap W_2$  has order  $q$  or  $q^2$ . In the first case stabilizer in  $S_0$  of  $t^g(W_1 \cap W_2)$  has order  $q^4$  and since  $t^g \cap t^g(W_1 \cap W_2) = t^g(C_{W_1 \cap W_2}(t^g))$ ,  $|C_{S_0}(t^g)| = q^3$ . But as  $C_{S_0}(t^g)$  contains no involution outside  $W_1 \cap W_2$ , this is impossible. In the second case  $W_1 \cap W_2 \sim V_i, i = 1$  or  $2$ , and  $C(V_i)$  contains a subgroup conjugate to  $C(W_1 \cap W_2) \geq S_0$ . This is impossible.

$$(3.5) \quad A/Z(A) \not\cong L_3(q).$$

*Proof.* Let  $G_0$  be as in (3.4). If  $d_1^{G_0} \cap S_0 = \emptyset$ , then by transfer  $S_0 \in \text{Syl}_2(\langle S_0^G \rangle)$ . McBride's result [12] gives a contradiction. Therefore  $d_1^g \in S_0$  for some  $g \in G_0$ . As  $A$  contains just one class of involutions,  $C_G(d_1)$  contains a conjugate  $S_0^g$  of  $S_0$  on which  $t$  acts. However  $d_1$  acts on  $A$  with  $C_A(d_1) \cong L_2(q)$  or  $U_3(q^{1/2})$ . In either case we have a contradiction by noting that  $C_A(d_1)$  is standard in  $C_G(d_1)$  and applying [6].

$$(3.6) \quad \bar{A} \not\cong L_n(q), n \geq 4.$$

*Proof.* As in (3.5) we may assume  $d_1^G \cap S_0 \neq \emptyset$ . Conjugating by an element of  $\langle N_1, A \rangle$  we may choose  $d_1^g \in S_0 - W_1$ . Say  $d_1^g$  has precisely  $l$  nontrivial Jordan blocks in its action on  $W_1$ , and choose a basis for  $W_1$  so that  $S_0/W_1$  is represented by lower triangular matrices. Then we may assume that  $d_1^g$  induces the matrix

$$\begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix}$$

on  $W_1$ , and we may write  $d_1^g = w_1v$  where  $v$  is an involution in  $V$ . Now  $W_1$



has the form  $W_1 = W_{11} \times W_{12}$ , where  $v$  centralizes  $W_{11}$ ,  $|W_{12}| = (q^2)^{2l}$ , and  $v$  induces  $l$  Jordan blocks on  $W_{12}$ . By (1.2) we may assume that  $w_1 \in W_{11}$ . Also we may choose  $W_{11}$  and  $W_{12}$  so that  $W_{11} \cap V_1 \neq 1$ . Then  $C_L(vW_1)$  contains a subgroup isomorphic to  $SL(n-2l, q)$  that centralizes  $W_{12}$  and is transitive on  $W_{11}^\#$ . Therefore we may assume that  $w_1 \in V_1 \cap W_{11}$ , so that  $d_1^q \in V \leq A$ . Now choose  $k$  minimal and a conjugate  $d_1^{q^{-1}}$  of  $d_1$  such that  $d_1^{q^{-1}} \in A$  and  $d_1^{q^{-1}}$  has precisely  $k$  nontrivial Jordan blocks on the usual  $\mathbb{F}_q$ -module for  $SL(n, q)$ .

The above arguments show that each involution in  $S_0$  is conjugate to one in  $A$ . Take  $d_1^q \in V$  and view  $S_0 \in \text{Syl}_2(SL(n, q^2))$ . We may assume that  $d_1^q$  is in Suzuki form ([2], §4). Let  $y \in S_0$  be any involution of Jordan rank  $r$  and in Suzuki form. Let  $M$  be the natural module for  $SL(n, q^2)$  and set  $J_r = C_{S_0}(C_M(y)) \cap C_{S_0}(M/[M, y])$ . Then  $J_r \triangleleft S_0$ ,  $C_M(J_r) = C_M(y)$ , and  $|J_r| = (q^2)^{r^2}$ .

Choose  $(S_0 \langle d_1 \rangle)^h$   $t$ -invariant with  $d_1 \in J_k^h \leq S_0^h$ . As in (3.3)  $S_0^h$  is characteristic in  $S_0^h \langle d_1^h, t \rangle$ . Then  $t$  acts on  $S_0^h$  and on a normal series of  $S_0^h$  with each factor a free  $\mathbb{F}_2 \langle t \rangle$ -module. Consequently each involution in  $S_0^h t$  is conjugate to  $t$ . In particular  $td_1 \sim t$ .

Suppose  $n$  is odd. Let  $X = (N_1 \cap N_2)'$ , so  $X$  induces  $SL(n-2, q^2)$  on  $W_1 W_2 / (W_1 \cap W_2)$ . By [2, (19.6)], each involution in  $Xtd_1$  is conjugate to an involution in  $W_1 W_2 td_1$ . Arguing as at the end of the second paragraph in the proof of (3.4), we have  $E(C(td_1)) \sim E(C(t))$  containing a section isomorphic to  $U_{n-2}(q)$  or  $Sp(n-3, q^2)$ . This is impossible.

Say  $n$  is even,  $n = 2b$ . Then  $J_b$  is the unique maximal elementary subgroup of  $S_0$  of order  $q^{2b}$ . Also  $J_k \leq I_b$ . So  $t$  normalizes  $J_b^h$ , centralizing a subgroup of order at least  $q^b$ . Thus  $C(t) \cap C(d_1)$  contains an elementary subgroup of order at least  $q^b$ . This is a contradiction and proves (3.6).

$$4. \widetilde{E(\bar{N}_i)} \cong L_{n-1}(q) \times L_{n-1}(q) \quad \text{FOR} \quad i = 1 \quad \text{AND} \quad 2$$

In this section we complete the proof of the main theorem by considering the case of (2.7) not treated in Section 3. That is, we assume that  $\widetilde{E(\bar{N}_i)} \cong L_{n-1}(q) \times L_{n-1}(q)$  for  $i = 1$  and  $2$ . Recall that  $G$  is a minimal counterexample to the main theorem.

We write  $\widetilde{E(\bar{N}_i)} = \widetilde{L_{i1}} \times \widetilde{L_{i2}}$ , where  $\bar{L}_{i1}$  and  $\bar{L}_{i2}$  are quasi-simple subgroups interchanged by  $t$ .

The first result shows that the module  $W_i$  decomposes under the action of  $\bar{L}_{i1} \bar{L}_{i2}$ .

(4.1)  $W_i = W_{i1} \times W_{i2}$  with  $\bar{L}_{ij} \cong SL(n-1, q)$  acting in the usual way on  $W_{ij}$  (when viewed as an  $\mathbb{F}_2$ -module) and centralizing  $W_{ik}$  for  $k \neq j$ .

*Proof.* We first claim that  $W_i$  when restricted to  $\bar{L}_{i1} \times \bar{L}_{i2}$  is reducible. Suppose this is not the case. Then  $W_i | \bar{L}_{i1}$  is homogeneous. If  $W_i | \bar{L}_{i1}$  were irreducible, then as  $\bar{L}_{i2} \leq C(\bar{L}_{i1})$ , Schur's lemma implies that  $\bar{L}_{i2}$  acts as a cyclic group on  $W_i$ ; impossible. Therefore  $W_i | \bar{L}_{i1}$  is reducible. If  $\bar{L}_{ij}$  acts nontrivially on an elementary Abelian 2-group of order  $2^a$ , then  $2^a \geq q^{n-1}$ . Indeed this follows from [3, (4B)] if  $n = 3$  and was proved in (3.1) if  $n \geq 4$ . So  $W_i | \bar{L}_{i1}$  is the sum of two  $\bar{L}_{i1}$ -invariant submodules  $Z_1, Z_2$  on which  $\bar{L}_{i1}$  induces equivalent representations.

Passing to the  $\mathbb{F}_q$ -module  $\hat{W}_i = W_i \otimes \mathbb{F}_q$  we have  $\hat{W}_i = M_1 \oplus \cdots \oplus M_r$ , where the  $M_i$  are absolutely irreducible, algebraically conjugate,  $\bar{L}_{i1}\bar{L}_{i2}$ -modules over  $\mathbb{F}_q$ . So each  $M_m = M_{m1} \otimes M_{m2}$  where  $M_{mj}$  is an (absolutely) irreducible  $\mathbb{F}_q$ -module for  $\bar{L}_{ij}$ . As an  $\mathbb{F}_q(\bar{L}_{i1})$ -module  $\hat{W}_i \cong (Z_1 \otimes \mathbb{F}_q) \oplus (Z_2 \otimes \mathbb{F}_q)$  and  $Z_1 \otimes \mathbb{F}_q \cong Z_2 \otimes \mathbb{F}_q$  is the sum of inequivalent, absolutely irreducible, modules for  $\bar{L}_{i1}$ . But  $M_m | \bar{L}_{i1}$  is the sum of  $\dim_{\mathbb{F}_q}(M_{m2})$  equivalent  $\mathbb{F}$ -modules for  $\bar{L}_{i1}$ . So as  $\dim(M_{m2}) \geq n - 1$ , we have  $n - 1 = 2$  and  $\bar{L}_{i1} \cong SL(2, q)$ .

It is easy to see that  $\bar{L}_{i1}$  stabilizes precisely  $q + 1$  subgroups of  $W_i$  of order  $q^2$ . So a Sylow 2-normalizer of  $\bar{L}_{i2}$ , say  $\bar{J}_{i2}$ , stabilizes one such subgroup  $I$  of  $W_i$ . As  $\bar{L}_{i1} \times \bar{J}_{i2}$  acts irreducibly on  $I$ ,  $O_2(\bar{J}_{i2})$  centralizes  $I$ . By symmetry the Sylow 2-subgroups of  $\bar{L}_{i1}$  centralize a subgroup of  $W_i$  of order at least  $q^2$ . It follows from [3(4B), (4C)] that  $\bar{L}_{i1}$  acts on  $I$  as on the usual module for  $\bar{L}_{i1} \cong SL(2, q)$  when viewed as an  $\mathbb{F}_2$ -module. From here it follows that each  $M_{mj}$  is an algebraic conjugate of the usual  $\mathbb{F}_q$ -module for  $\bar{L}_{ij}$ . However, we know also that  $L_i$  acts on  $W_i/V_i$  and on  $V_i$  as on the usual module for  $SL(2, q)$ , viewed as an  $\mathbb{F}_2$ -module. So if we let  $h$  be an element of order  $q - 1$  in  $L_i$ , then there is an element  $\alpha \in \mathbb{F}_q^\#$  such that the eigenvectors of  $h$  on  $\hat{W}_i$  are the algebraic conjugates of  $\alpha$  and  $\alpha^{-1}$ , each with multiplicity two. This is not consistent with the form of  $M_1, \dots, M_r$ . Therefore  $W_i$  is reducible when restricted to  $\bar{L}_{i1}\bar{L}_{i2}$ , as claimed.

Let  $W_{i1}$  be an irreducible submodule of  $W_i$  under  $\bar{L}_{i1}\bar{L}_{i2}$ , and choose notation so that  $\bar{L}_{i1}$  is not trivial on  $W_{i1}$ . Then as before  $|W_{i1}| \geq q^{n-1}$ , and since  $W_i/W_{i1}$  is also nontrivial  $|W_{i1}| = |W_i/W_{i1}| = q^{n-1}$ . Moreover  $\bar{L}_{i1}$  acts irreducibly on  $W_{i1}$ . Then  $\bar{L}_{i2} \leq C(\bar{L}_{i1})$  forces  $\bar{L}_{i2}$  to be trivial on  $W_{i1}$ . As  $t$  interchanges  $\bar{L}_{i1}$  and  $\bar{L}_{i2}$  we set  $W_{i2} = W_{i1}^t$  and get  $W_i = W_{i1} \times W_{i2}$ .

It remains to show that as an  $\mathbb{F}_2$ -module,  $W_{ij}$  is equivalent to the usual module for  $\bar{L}_{ij}$ . But this follows from the fact that  $L_i$  acts on  $W_i/V_i$  and on  $V_i$  as on the usual module. The proof of (4.1) is complete.

(4.2) *Notation.* Let  $S_0^t \in \text{Syl}_2(N_1^{(\infty)})$  with  $C_{S_0}(t) = V$ . By (4.1) we have  $S_0 = A_1 \times B_1$  with  $B_1 = A_1^t$ . Here  $A_1 \geq W_{11}$  and  $B_1 \geq W_{12}$ . As  $V \in \text{Syl}_2(L_n(q))$  this is also true of  $A_1$  and  $B_1$ .

Recall the  $(B, N)$ -notation for  $A$ . We have a root system  $\Sigma$  and  $V = \prod_{r \in \Sigma^+} U_r = \langle U_{\alpha_1}, \dots, U_{\alpha_{n-1}} \rangle$ . Accordingly we write

$$A_1 = \prod_{r \in \Sigma^+} Y_r, B_1 = \prod_{r \in \Sigma^+} Z_r, \text{ where } U_r \leq Y_r \times Z_r \text{ and } Z_r = Y_r^t.$$

Let the Weyl group of  $A$  be generated by reflections  $s_1, \dots, s_{n-1}$  with  $s_i \in \langle U_{\alpha_i}, U_{-\alpha_i} \rangle$  and  $U_{-\alpha_i} = U_{\alpha_i}^{s_i}$ .

(4.3) *There exist subgroups  $G_1, G_2$  of  $G$  with the following properties:*

- (i)  $G_1^t = G_2$ .
- (ii)  $[G_1, G_2] = 1$ .
- (iii)  $G_1 \cap G_2 \leq Z(G_1 G_2)$  and  $|Z(G_1 G_2)|$  is odd.
- (iv)  $G_1 G_2 / Z(G_1 G_2) \cong L_n(q) \times L_n(q)$ .

*Proof.* First note that in the group  $N_2^{(\infty)}$  we have subgroups corresponding to those in (4.2) which we label  $\hat{A}_1, \hat{B}_1, \hat{Y}_r, \hat{W}_{21}, \hat{W}_{22}$ . Then  $V = C(t) \cap (\hat{A}_1 \times \hat{B}_1)$ . We know that  $C(V_2) \cap A_1 B_1$  is elementary of order  $(q^{n-1})^2 = |W_2|$ , so we may take  $W_2 \leq A_1 B_1$ . It then follows that  $W_2 \triangleleft A_1 B_1$  and conjugating if necessary we may assume  $A_1 B_1 = \hat{A}_1 \hat{B}_1 = S_0$ .

We have  $N_1^{(\infty)} = M_{11} \times M_{12}$  where  $W_{1i} = O_2(M_{1i})$  and  $\bar{M}_{1i} = \bar{L}_{1i}$ . As  $t$  interchanges  $M_{11}$  and  $M_{12}$ , and  $C_{N^{(\infty)}}(t) = V_1 L_1$ , we may write  $M_{1i} = W_{1i} L_{1i}$ , where  $L_{1i} \cong SL(n-1, q)$ ,  $L_{12} = L_{11}^t$ , and  $L_1 = C(t) \cap L_{11} L_{12}$ . We may assume that  $W_{11} = Y_{\alpha_1} \dots Y_{\alpha_1 + \dots + \alpha_{n-1}}$  and  $A_1 \cap L_{11} = \langle Y_{\alpha_2}, \dots, Y_{\alpha_{n-1}} \rangle$ . Then  $B_1 \cap L_{12} = \langle Z_{\alpha_2}, \dots, Z_{\alpha_{n-1}} \rangle$ . Also  $\bar{L}_{11} = \langle Y_{\pm \alpha_2}, \dots, Y_{\pm \alpha_{n-1}} \rangle$ , where for  $i = 2, \dots, n-1$  we write  $Y_{-\alpha_i} = Y_{\alpha_i}^{s_i}$ . Similarly  $L_{12} = \langle Z_{\pm \alpha_2}, \dots, Z_{\pm \alpha_{n-1}} \rangle$ .

The Krull-Schmidt theorem implies that with a possible change of notation  $\hat{A}_1 \leq A_1 Z(S_0)$  and  $\hat{B}_1 \leq B_1 Z(S_0)$ . We claim that  $A_1 = \hat{A}_1$  and  $B_1 = \hat{B}_1$ . Suppose that  $n = 3$ . Then  $q \geq 4$  and  $N_1 \cap N(W_1 W_2)$  contains an Abelian subgroup  $X_1 X_2$  such that  $L_{1i} \geq X_i$  is cyclic of order  $q-1$  and fixed-point-free on  $S_0 \cap M_{1i}$ . Viewing this in  $N_2$  we conclude that  $A_1 = \hat{A}_1$  and  $B_1 = \hat{B}_1$ .

Now suppose  $n \geq 4$ . Then  $L_{11}$  contains a subgroup  $X_1 \cong SL(n-2, q)$  normalizing  $W_2$ . Namely,  $X_1 = \langle Y_{\pm \alpha_2}, \dots, Y_{\pm \alpha_{n-2}} \rangle$ . Then  $[X_1, W_1 W_2] = W_1 W_2 \cap M_{11}$ . Considering  $X_1 \times X_1^t \leq N_2^{(\infty)}$  we see that  $X_1$  induces a subgroup of  $\bar{L}_{21}$  on  $W_1 W_2$ . Therefore  $[X_1, W_1 W_2] \leq M_{21}$ . Therefore  $W_1 W_2 \cap M_{11} \leq \hat{A}_1$ . Since  $A_1 \cap X_1 \leq \hat{A}_1$  we have  $A_1 = (A_1 \cap X_1)(W_1 W_2 \cap M_{11}) \leq \hat{A}_1$ , so by orders  $A_1 = \hat{A}_1$ . Similarly  $B_1 = \hat{B}_1$ .

Now write  $N_2^{(\infty)} = \bar{M}_{21} \times \bar{M}_{22}$ , where  $\bar{M}_{2i} = \bar{W}_{2i} \bar{L}_{2i}$ . Then  $\bar{W}_{21} = Y_{\alpha_{n-1}} \dots Y_{\alpha_1 + \dots + \alpha_{n-1}}$ ,  $\bar{W}_{22} = Z_{\alpha_{n-1}} \dots Z_{\alpha_1 + \dots + \alpha_{n-1}}$ ,  $A_1 \in \text{Syl}_2(\bar{M}_{21})$ , and  $B_1 \in \text{Syl}_2(\bar{M}_{22})$ . We may also choose  $\bar{L}_{21}, \bar{L}_{22}$  to be interchanged by  $t$  and  $\langle U_{\pm \alpha_1}, \dots, U_{\pm \alpha_{n-2}} \rangle = L_2 = C(t) \cap (L_{21} L_{22})$ . Then  $\bar{L}_{21}$  can be expressed as  $\bar{L}_{21} = \langle \hat{Y}_{\pm \alpha_1}, \dots, \hat{Y}_{\pm \alpha_{n-2}} \rangle$  and  $\bar{L}_{22} = \langle \hat{Z}_{\pm \alpha_1}, \dots, \hat{Z}_{\pm \alpha_{n-2}} \rangle$ . We know that  $A_1 \leq \bar{M}_{21}$  and  $C_G(t) \cap \bar{M}_{21} = 1$ . Considering the map  $g \rightarrow [g, t]$  for  $g \in M_{21}$ , we get  $\hat{Y}_{\alpha_i} = Y_{\alpha_i}$  for  $i = 1, \dots, n-2$ .

Let  $w_0 \in \langle s_1, \dots, s_{n-1} \rangle$  be the word of greatest length in the generators  $s_1, \dots, s_{n-1}$ . Fix  $i \in \{2, \dots, n-1\}$ . Then  $\alpha_i^{w_0} = -\alpha_{n-i}$ ;  $w_0$  can be expressed as  $w_0 = s_i s_{j_1} \dots s_{j_r}$ , where  $r+1 = l(w_0)$ , and for  $1 \leq k \leq r$ ,  $(\alpha_i) s_{j_1}, \dots, s_{j_k} =$

$(-\alpha_i) s_i s_{j_1}, \dots, s_{j_k} \in \Sigma^+$ . As each  $j_k \in L_1$  or  $L_2$ , we conclude that  $(Y_{-\alpha_i})^{w_0} = (Y_{\alpha_i})^{s_{j_1} \dots s_{j_r}} = Y_{\alpha_{n-i}}$ . Also for  $i = \{1, \dots, n-2\}$  we have

$$(\hat{Y}_{-\alpha_i})^{w_0} = (\hat{Y}_{-\alpha_i})^{s_i s_{j_1} \dots s_{j_r}} = (Y_{\alpha_i})^{s_{j_1} \dots s_{j_r}} = Y_{\alpha_{n-i}}.$$

If  $i \geq 2$ , then  $(\hat{Y}_{-\alpha_i})^{w_0} = (Y_{-\alpha_i})^{w_0}$  so  $Y_{-\alpha_i} = \hat{Y}_{-\alpha_i}$ . Now set  $Y_{-\alpha_1} = \hat{Y}_{-\alpha_1}$ . Conjugating these equations by  $w_0$  and making similar arguments for the root groups  $Z_{\pm\alpha_i}$ , we have

$$(Y_{\pm\alpha_i})^{w_0} = Y_{\mp\alpha_{n-i}} \quad \text{and} \quad (Z_{\pm\alpha_i})^{w_0} = Z_{\mp\alpha_{n-i}} \quad \text{for } i = \{1, \dots, n-1\}.$$

Notice that  $G_1 = \langle L_{11}, L_{21} \rangle = \langle Y_{\pm\alpha_1}, \dots, Y_{\pm\alpha_{n-1}} \rangle \leq C(B_1)$ . By the above,  $w_0$  normalizes  $\langle L_{11}, L_{21} \rangle$  and  $\langle B_1, B_1^{w_0} \rangle = \langle Z_{\pm\alpha_1}, \dots, Z_{\pm\alpha_{n-1}} \rangle = G_2$ . Therefore  $[G_1, G_2] = 1$  and  $G_2 = G_1^t$ , proving the first two parts of (4.3).

Clearly  $G_1 \cap G_2 \leq Z(G_1 G_2) = Z$  and  $A \leq G_1 G_2$ . As  $Z(G_1 G_2) \leq C_G(A)$ ,  $|Z(G_1 G_2)|$  is odd, proving (iii). This also implies that  $C(t)$  covers  $C(t) \cap \widetilde{G_1 G_2} = C(t) \cap (G_1 Z/Z) \times (G_2 Z/Z)$ .  $G_1$  is generated by the perfect groups  $L_{11}$  and  $L_{21}$ , so  $G_1$ , and therefore  $G_2$ , is perfect. Consequently  $G_1 G_2 / Z(G_1 G_2) \cong A/Z(A) \times A/Z(A) \cong L_n(q) \times L_n(q)$ , completing the proof of (4.3).

All that remains is to prove that  $G_0 = G_1 G_2 \trianglelefteq G$ . For this set  $\Omega = \{G_0^g : g \in G\}$ .

(4.4)  $t$  fixes a unique point of  $\Omega$ , so  $C(t) \leq N(G_0)$ .

*Proof.* Suppose  $t$  fixes  $G_0^g$ . If  $t \in N(G_1^g) \cap N(G_2^g)$ , then checking  $C_{G_0^g}(t)$  we get a contradiction. So  $(G_1^g)^t = G_2^g$  and hence  $E(C_{G_0^g}(t))$  involves  $L_n(q)$ . We conclude that  $A \leq G_0^g$ . Arguing as in the proof of (2.7) we see that  $N_G(V_i)^\infty$  centralizes  $O(N_G(V_i))$  and hence  $W_i = O_2(N_G(V_i))$ . So  $W_i$  is the unique Sylow 2-subgroup of  $C_G(V_i)$ . As  $(C_{G_0^g}(V_i))^{(\infty)}$  has the form  $D_i L_i$ , where  $D_i$  is normal and elementary of order  $|V_i|^2$ ,  $D_i = W_i$ . Also  $G_0^g = \langle N_{G_0^g}(D_1)^{(\infty)}, N_{G_0^g}(D_2)^{(\infty)} \rangle = \langle N_{G_0^g}(W_1)^{(\infty)}, N_{G_0^g}(W_2)^{(\infty)} \rangle$  and  $N_{G_0^g}(W_i)^{(\infty)} \cong N_G(W_i)^{(\infty)}$ . But  $G_0 = \langle N(W_1)^{(\infty)}, N(W_2)^{(\infty)} \rangle$ , so  $G_0 = G_0^g$ , proving the result.

(4.5)  $G$  contains a normal subgroup  $G_0$  with  $G = G_0 \langle t \rangle$  and  $t \notin G_0$ .

*Proof.* Let  $X = N_G(G_1) \cap N_G(G_2) \leq N_G(G_0)$ . Then  $X \langle t \rangle$  contains a Sylow 2-subgroup of  $G$ . By the Thompson transfer lemma we need only show that  $t^G \cap X = \emptyset$ . Suppose  $t^g \in X$ . Then  $t \in X^{g^{-1}} \leq N_G(G_0^{g^{-1}})$ , so by (4.4),  $g^{-1} \in N(G_0)$ . But as  $X \trianglelefteq N_G(G_0)$  this is impossible.

(4.6) Let  $g \in \text{Aut}(S_1)$  have odd order, where  $S_1 \in \text{Syl}_2(G_1)$ . Then  $g$  normalizes  $W_{11}$  and  $W_{21}$ .

*Proof.* We have  $S_1 \in \text{Syl}_2(L_n(q))$ , so if  $q \geq 4$  the result follows from [11]. Suppose  $q = 2$ . Let  $X = W_{11}$  and  $Y = W_{11}^g$ . Then  $X$ , and hence  $Y$ , has the property that  $(X \cap Z_i(S_1))/(X \cap Z_{i-1}(S_1))$  has order 2 for each  $i = 1, \dots, n-1$ ,

where  $Z_i(S_1)$  is the  $i$ th term of the upper central series of  $S_1$ . As  $g$  normalizes  $Z(S_1)$ ,  $X \cap Y \geq Z(S_1)$ . Choose  $i$  maximal with  $X \cap Z_i(S_1) = Y \cap Z_i(S_1)$ .

First suppose  $i \geq 2$ . If  $i = n - 1$ , then  $X = Y$  and we have  $g$  normalizing  $W_{11}$ . Each factor  $Z_{i+1}(S_1)/Z_i(S_1)$  is the direct product of images of certain root subgroups of  $S_1$ . For example  $Z_{n-2}(S_1)/Z_{n-3}(S_1) \cong Y_{\alpha_1+\alpha_2} \times Y_{\alpha_2+\alpha_3} \times \cdots \times Y_{\alpha_{n-2}+\alpha_{n-1}}$ . Using the commutator relations and the fact that  $[Y \cap Z_{i+1}(S_1), S_1] \leq Y \cap Z_i(S_1) = X \cap Z_i(S_1)$  we see that  $(Y \cap Z_{i+1}(S_1)) Z_i(S_1) = (X \cap Z_{i+1}(S_1)) Z_i(S_1)$ . Let  $y \in Y \cap (Z_{i+1}(S_1) - Z_i(S_1))$  and write  $y = xv$  where  $x \in X \cap Z_{i+1}(S_1)$  and  $v \in Z_i(S_1)$ . If  $v \notin (X \cap Z_i(S_1)) Z_2(S_1)$ , then this leads to a contradiction. This is because it is then possible to choose an element  $s \in S_1$  such that  $[x, s] \in X \cap Z_i(S_1)$  and  $[v, s] \notin X \cap Z_i(S_1)$ . So  $v \in (X \cap Z_i(S_1)) Z_2(S_1)$ . Now  $g$  normalizes each  $X \cap Z_j(S_1) = Y \cap Z_j(S_1)$  for  $j \leq i$ , and also  $g$  normalizes  $Z_2(S_1)$  which has order 8. It follows that  $g$  stabilizes a series of  $(X \cap Z_{i+1}(S_1)) Z_2(S_1)$  each factor of which has order 2. Since  $|g|$  is odd,  $g$  must centralize  $(X \cap Z_{i+1}(S_1)) Z_2(S_1)$ . In particular  $g$  centralizes  $X \cap Z_{i+1}(S_1)$ , which is not the case.

Therefore  $i = 1$ . In particular this implies  $X \cap Y = Z(S_1)$ , so  $[X, Y] \leq Z(S_1)$  and  $Y \leq C(X/Z(S_1)) = W_{11}W_{21}$ . Then  $W_{11}W_{21} = XY = C_{S_1}(X/Z(S_1)) = C_{S_1}(Y/Z(S_1))$  is  $g$ -invariant. Now  $S_1/XY$  is isomorphic to the Sylow 2-subgroups of  $SL(n-2, 2)$ . We may apply induction unless  $n-2 = 2, 3$ , or  $4$ . Note that  $n-2 = 2$  does not occur as  $\tilde{A} \not\cong L_4(2)$ . Assume for the moment that  $n-2 > 4$ .

Inductively  $g$  normalizes subgroups  $H/XY, K/XY$ , with  $H/XY, K/XY$  analogous to  $W_{11}, W_{21}$ , respectively. Relabeling, if necessary, we have the centralizer in  $W_{i1}Z(XY)/Z(XY)$  of  $H/Z(XY)$ , respectively, a 1-space or hyperplane of  $W_{i1}Z(XY)/Z(XY)$ . For  $K/Z(XY)$  the centralizers are, respectively, a hyperplane or 1-space of  $W_{i1}Z(XY)/Z(XY)$ . Now  $g$  normalizes  $C(K/Z(XY)) \cap XY/Z(XY)$ . This forces  $C(H/XY) \cap X/Z(XY) \cap X^g/Z(XY) \neq 1$ . However  $X \cap X^g = X \cap Y = Z(XY)$ , a contradiction.

We still have the cases  $n = 5, 6$ . If  $n = 5$  we have  $S/XY \cong D_8$  so  $g$  centralizes  $S/XY$  and the above argument works. Let  $n = 6$ . If  $g$  normalizes  $H/XY, K/XY$  we argue as before. Suppose  $g$  does not normalize  $H/XY, K/XY$ . Argue as in the second paragraph of the proof to conclude that  $H/XY \cap K/XY = Z(S_1/XY)$ . Then  $\langle g \rangle$  must induce  $Z_3$  on  $Z_2(S_1/XY)$ . So  $\langle g \rangle$  is transitive on the three Klein groups of  $Z_2(S_1/XY)$  containing  $Z(S_1/XY)$ . Considering the orders of the centralizers of these subgroups in their action on  $XY/Z(XY)$  we obtain a contradiction. We have now shown that  $W_{11}^g = W_{11}$ . Similarly  $W_{21}^g = W_{21}$ .

At this point we can complete the proof of the main theorem. We assume condition (\*) holds. It suffices to show  $G_1G_2 \trianglelefteq G$  and this follows from [16, (2.7)] once the hypotheses of that result are verified. For this we use [16, (2.6)]. Suppose  $G_1G_2 \not\trianglelefteq G$ . We conclude that there exists a conjugate  $t_1$  of  $t$  such that  $t_1 \notin N(G_1G_2)$  and  $tt_1 \in N(S_i)$ ,  $i = 1, 2$ . As in the proof of [16, (2.2)],  $O_2(\langle t, t_1 \rangle) \leq N(G_1G_2)$ . So we may assume that  $g = tt_1$  has odd order. By (4.6)  $g$

normalizes  $W_1$  and  $W_2$ . However,  $G_0 = G_1G_2 = \langle N_G(W_1)^{(\infty)}, N_G(W_2)^{(\infty)} \rangle$ , so  $G_0 = G_0^g$ . This contradiction completes the proof of the main theorem.

## REFERENCES

1. M. ASCHBACHER, Standard component of alternating type centralized by a 4-group, to appear.
2. M. ASCHBACHER AND G. SEITZ, Involutions in Chevalley groups over fields of even order, *Nagoya Math. J.* **63** (1976), 1-91.
3. P. FONG AND G. SEITZ, Groups with a  $(B, N)$ -pair of rank 2, I, *Invent. Math.* **21** (1973), 1-57.
4. D. GOLDSCHMIDT, 2-fusion in finite groups, *Ann. Math. (2)* **99** (1974), 70-117.
5. R. GRIESS, "Schur Multipliers of the Known Simple Groups," Thesis, University of Chicago, 1971.
6. R. GRIESS, D. MASON, AND G. SEITZ, Bender groups as standard component, to appear.
7. K. HARADA, On finite groups having self-centralizing 2-subgroups of small order, *J. Algebra* **33** (1975), 144-160.
8. G. HIGMAN, Suzuki 2-groups, *Illinois J. Math.* **7** (1963), 79-96.
9. B. HUPPERT, "Endliche Gruppen, I," Springer-Verlag, Berlin/Heidelberg/New York, 1960.
10. S. LANG, Algebraic groups over finite fields, *Amer. J. Math.* **18** (1956), 555-563.
11. P. MCBRIDE, Automorphisms of 2-groups, to appear.
12. P. MCBRIDE, "A Classification of Groups of Type  $A_n(q)$ , for  $n \geq 3$  and  $q = 2^k \geq 4$ ," Ph.D. Thesis, University of Illinois at Chicago Circle, 1975.
13. M. O'NAN, A characterization of  $L_n(q)$  as a permutation group, *Math. Z.* **127** (1972), 301-314.
14. F. PIPER, On elations of finite projective spaces, *Geom. Ded.* **2** (1973), 13-27.
15. J. OSTROM AND A. WAGNER, On projective and affine planes with transitive collineation groups, *Math. Z.* **71** (1959), 186-199.
16. G. SEITZ, Chevalley groups as standard components, I, II, III.
17. E. SHULT, On the fusion of an involution in its centralizer, to appear.
18. R. SOLOMON, Standard components of alternating type, to appear.
19. R. STEINBERG, Representations of algebraic groups, *Nagoya Math. J.* **22** (1963), 33-56.
20. J. THOMPSON, Fixed points of  $p$ -groups acting on  $p$ -groups, *Math. Z.* **86** (1964), 12-13.
21. M. ASCHBACHER AND G. SEITZ, On groups with a standard component of known type, *Osaka J. Math.* **13** (1976), 439-482.
22. G. SEITZ, Small rank permutation representations of finite Chevalley groups, *J. Algebra* **28** (1974), 508-517.
23. R. STEINBERG, Générateurs, relations, et, revêtements de groupes algébriques, colloque sur la théorie des groupes algébriques, Brussels, 1962, pp. 113-127.